

## REFERENCES

- Abbott, S. A. (1998). The Benefits of Patient Education. *Gastroenterology Nursing*, 21(5), 207-209.
- Agarwal, R. (2000). Assessment of Blood Pressure in Hemodialysis patients. *Seminars in Dialysis*, 15(5), 299-304.
- Agarwal, R. (2005). Hypertension and survival in chronic hemodialysis patients—Past lessons and future opportunities. *Kidney International*, 67, 1–13.
- Agarwal, R. (2010). Blood Pressure and mortality among hemodialysis patients. *Hypertension*, 55, 762-768.
- Agarwal, R. (2010). Hypervolemia is associated with increased mortality among hemodialysis patients. *Hypertension*, 56,512-517.
- Agarwal, R. (2012). How can we prevent intradialytic hypotension? *Current Opinion in Nephrology and Hypertension*, 21(6), 593–599.
- Agarwal, R., & Weir, M.R. (2010). Dry-Weight: A Concept Revisited in an Effort to Avoid Medication-Directed Approaches for Blood Pressure Control in Hemodialysis Patients. *Journal of the American Society of Nephrology*, 5(7): 1255–1260.
- Ash, S., Campbell, K., MacLaughlin, H., McCoy, E., Chan, M., Anderson, K., et al. (2006). Evidence based practice guidelines for the nutritional management of chronic kidney disease. *Nutrition & Dietetics*, 63, S33-S45.
- Bander, S.L., & Walters, B.A.J. (1998). Hemodialysis morbidity and mortality: links to patient non-compliance. *Current Opinion in Nephrology and Hypertension*, 7(6), 649-653.
- Bandura, A. (1977). Toward a unifying theory of behavioral change psychological review, 84, 191-215.

- Baraz, S., Parvardeh, S., Mohammadi, E., & Broumand, B. (2010). Dietary and fluid compliance: an educational intervention for patients having haemodialysis. *Journal of Advanced Nursing*, 66, 60-68.
- Barnett, T., Tang, L. Y., Pinikahana, J., & Tan, S.Y. (2008). Fluid compliance among patients having haemodialysis: can an educational programme make a difference? *Journal of Advanced Nursing*, 61(3), 300–306.
- Batson, H., & Schwartz, M. (2000). Effective technique to increase hemodialysis and PD patient compliance, Part II. *Nephrology News Issues*, 13(6), 39-40, 42.
- Bittenbinder, E.N. (2008). The Role of Patient Education in the Patient's Familiarity and Understanding of treatment, Internship practicum report. *Proquest Dissertations & Thesis, University of North Texas, Health Science Center*.
- Bland R.J., Cottrell, R.R., & Guyler, L.R. (2008). Medication compliance of hemodialysis patients and factors contributing to non-compliance. *Dialysis & Transplantation* 37, 174–178.
- Bleyer, A.J., Russel, G.B., & Satko, S.G. (1999). Sudden and cardiac death rates in hemodialysis patients. *Kidney International*, 55(4), 1553-1559.
- Burge, S., White, D., Bajorek, E., Bazaldua, O., Trevino, J., Albright, T., et al. (2005). Correlates of Medication Knowledge and Adherence. *Family Medicine*, 37(10):712-718.
- Casey, J., Johnson, V., & McClelland, P. (2002). Impact of stepped verbal and written reinforcement of fluid balance advice within an outpatient haemodialysis unit: a pilot study. *Journal of Human Nutrition and Dietetics*, 15(1), 43-47.
- Chalmers, J., MacMahon, S., & Manica, G. (1999). World health Organization-International Society of Hypertension Guidelines for the management of hypertension. Guidelines sub-committee of the World Health Organization. *Clinical Experience Hypertension*, 21, 1009-1060.

Chan, Y.M., Zalilah, M.S., Hii, S.Z. (2012). Determinants of Compliance behaviors among patients Undergoing Hemodialysis in Malaysia (Electronic version). *PLoS ONE*, 7(8), e41362. doi:10.1371/journal.pone.0041362.

Chamney, M, Pugh-Clarke, K., & Kafkia, T. (2009). Management of co-morbid diseases in a patient with established renal failure. *Journal of Renal Care* 35, 151-158.

Charra, B. (2007). Fluid balance, dry weight, and blood pressure in dialysis. *Hemodialysis International*, 11, 21-31.

Chen, C.H., Lin, Y.P., Yang, W.C., & Ding, Y.A. (2003). Volume status and Blood Pressure during Long Term Hemodialysis: Role of Ventricular Stiffness. *Hypertension*, 42(3), 257-262.

Christensen, A. J., Moran, P. J., Wiebe, J. S., Ehlers, S. L., & Lawton, W. J. (2002). Effect of behavioral self-regulation interventions on patient adherence in hemodialysis. *Health Psychology*, 21(4), 393 – 397.

Chorold, L.B., & Lanchester, L.E. (1995). *ANNA, Core Curriculum for Nephrology Nursing* (3rd ed.). New Jersey, Pitman: American Nephrology Nurses' Association.

Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences* (2 nd .ed). Hilldale, Lawrence Erlbaum Associates.

Couchoud, C., Kooman, J., Finne, P., Leivestad, T., Stojceva-Taneva, O., Ponikvar, J.B., et al. (2009). From registry data collection to international comparisons: examples of haemodialysis duration and frequency. *Nephrology Dialysis Transplant*, 24, 217-224.

Cvengros, J. A., Christensen, A. J., & Lawton, W. J. (2004). The role of perceived control and preference for control in adherence to chronic medical regimen. *Annals of Behavior Medicine*, 27(3), 155 – 161.

Denhaerynck, K., Manhaeve, D., Dobbels, F., Garzoni, D., Nolte, C., Geets, D.S. (2007) Prevalence and consequences of non-adherences to hemodialysis regimes. *American Journal of Critical Care*, 16, 222-235.

Fincham,D.,Kagee,A.,&Moosa,M.R.(2008). Dietary and fluid adherence among haemodialysis patients attending public sector hospitals in the Western Cape. *South Africa Journal of Clinical Nutrition*, 21(2),7-19.

Fishbane, S.A., & Scribner, B.H. (2002). Blood Pressure Control in Dialysis Patients. *Seminars in Dialysis*, 15(3), 144-145.

Fisher,L.,Cairns,H.S.,Amir-Asari,B.,Scoble,J.E.,Chalder,T.G.,& Tressure,J.(2006). Psychological intervention in fluid management. *Palliative Support Care*, 41, 419 - 424.

Friedman, E.A. (2001). Must we treat Noncompliant ESRD Patients? *Seminars In Dialysis*, 14(1), 23-27.

Forbes, A., & While, A. (2009). The nursing contribution to chronic disease management: a discussion paper. *International Journal of Nursing Studies*, 46, 119-130.

Golper, T. (2001). Patient education: can it maximize the success of therapy? *Nephrology Dialysis Transplantation*, 16(suppl 7), 20-24.

Goovaerts,T., Jadoul,M., & Goffin,E.(2005).Influence of a Pre-Dialysis Education Programme (PDEP) on the mode of renal replacement therapy. *Nephrology Dialysis Transplantation*, 20, 1842–1847.

Horl, M.P., & Horl, W.H. (2002).Hemodialysis-associated hypertension: Pathophysiology and therapy. *American Journal of Kidney Disease*, 39:227–244.

Ifudu, O., Uribarri, J., Rajwani, I., Vlacich, V., Reydel, K., Delosreyes, G., & Friedman, E.A. (2002). Relation between interdialytic weight gain, body weight and nutrition in hemodialysis patients. *American Journal of Nephrology*, 22, 363-368.

Inrig, J.K., Uptal, D.P., Gillespie,B.S., Hasselblad,V., Himmelfarb,J., Reddan,D., et al. (2007). Relationship between Interdialytic Weight Gain and Blood Pressure Among Prevalent Hemodialysis Patients. *American Journal of Kidney Disease*. 50 (1):108–118.

Julius,J.R., Novitsky,J.R.,M.A.,& Dubin,W.R.(2009).Medication adherence :A review of the Literature and implication for clinical practice. *Journal of Psychiatric Practice*, 15(1), 34-44.

KalantarZadeh,K.,Regodir,D.L.,Kovesdhy,C.P.,Wyck,D.,Bunnapradist,S.,Horwich,T.B, Fonarow,C.G.(2009). Fluid retention is associated with cardiovascular mortality in patients undergoing Long –term hemodialysis.*Circulation*,119, 671-679.

Klassen, P.S., Lowrie, E.G., & Reddan, D.N. (2002). Association between pulse pressure and mortality in patients undergoing maintenance haemodialysis, *JAMA*, 287, 1548-1555.

Karakurt, P., & Kasıkçı, M.K. (2012). The effect of education given to patients with Type 2 Diabetes Mellitus on self-care. *International Journal of Nursing Practice*, 18, 170–179.

Karamanidou, C., Weinman, J., & Horne, R.(2007). Review of the prevalence and determinants of adherence to phosphate-binding medication in renal dialysis patients. *Nephrology Dialysis Transplantation* 22, 375.

Kauric-Klein, Z., & Artinian, N. (2007). Improving blood pressure control in hypertensive hemodialysis patients. *The CANNT Journal*, 17(4), 24 -29.

Kessels, R.P.C.(2003) Patients' memory for medical information *Journal of the Royal Society of Medicine*,96, 219-222.

Khechane, N. L. & Mwaba, K. (2004). Treatment adherence and coping with stress among black South African haemodialysis patients. *Social Behavior and Personality*, 32(8), 777 – 782.

Kimmel, P.L., Peterson, R.A., Weihs, K.L., Simmens, S.J., Alleyne, S., Cruz, I., & Veis, J.H. (1998). Psychosocial factors, behavioral compliance and survival in urban hemodialysis patients. *Kidney International*, 54(1), 245-254.

Kimmel, P.L., Varela, M.P., Peterson, R.A., Weihs, K.L., Simments, S.J., Alleyne, S., Amarashine, A., Mishkin, G.J., Cruz, I., & Veis, J.H. (2000). Interdialytic weight gain and survival in hemodialysis patients: Effects of duration of ESRD and diabetic mellitus. *Kidney International*, 57(3), 1141-1151.

Kugler, C., Vlaminck, H., Haverich, A. & Maes, B. (2005): Nonadherence with diet and fluid restrictions among adults having hemodialysis. *Journal of Nursing Scholarship*, 37, 25-29.

Kutner, N.G. (2001). Improving Compliance in Dialysis Patients: Does Anything Work? *Seminars in Dialysis*, 14(5), 324-327.

Kyngas, H., Duffy, M., Kroll, T. (2000). Conceptual analysis of compliance. *Journal of Clinical Nursing*, 9(1), 5-12.

Lee, A., Song, X., Khan, I., Belozeroff, V., Goodman, W., Fulcher, N., Diakun, D.(2011). Association of cinacalcet adherence and costs in patients on dialysis. *Jounal of Medical Economic*,14(6),798-804.

Leunissen,K.M.,Kooman,J.P.,& Luik,A.J.(1999).How important is volume excess in the etiology of hypertension in dialysis patients ?(Opinion).*Seminars in Dialysis*, 12(5),299-301.

Levey, A.S., Eckardt. K.U., Tsukamoto. Y., Levin. A., Coresh, J., Rossert. J., De Zeeuw, D., Hostetter, T.H., Lameire, N. & Eknayan, G. (2005). Definition and classification of chronic kidney disease: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney International*, 67, 2089-2100.

Leypoldt, J.K., Cheung, A.K., Delmez, J.A., Gassman, J.J., Levin, N.W., Lewis, J.A.B., Lewis, J.L., & Rocco, M.V. (2002). Relationship between volume status and blood pressure during chronic hemodialysis. *Kidney International*, 61(1), 266-275.

Lindberg, M., Prutz, K-G., Linberg,P.,& Wikstrom,B.(2009). Interdialytic weight gain and ultrafiltration rate in hemodialysis: Lessons about fluid adherence from a national registry of clinical practice. *Hemodialysis International*,13,181-188.

Lindberg, M.(2010). Excessive Fluid Overload Among Haemodialysis Patietns.Prevalence, Individual Characteristics and Self-regulation of fluid intake. *Digital Comprehensive Summaries of Uppsala Dissertations from the Faculty of Medicine 551*, Acta Universitatis Upsaliensis.

Luik, A.J.,Weideman, P., Cheriex, E., Kooman, J.P., Leunissen, K.M. (2001).The influence of increasing dialysis treatment time and reducing dry weight on blood pressure control in hemodialysis patients: a prospective study. *American Journal of Nephrology*,21(6),471-478.

Locatelli, F., Marcelli, D., & Conte, F. (2000). Cardiovascular disease in chronic renal failure: the challenge continues. *Nephrology Dialysis Transplantation*, 15(suppl 3), 69-80.

Lopez-Gomez, J.M., Villaverde, M., Joffre, R., Rodriguez-benitez, P., & Perez-Garcia, R. (2005). Interdialytic weight gain as blood marker of blood pressure, nutrition, and survival in hemodialysis patients. *Kidney International*, 67(suppl 93), 63-68.

Lucas, M.F., Quereda, C., & Teruel, J.L. (2003). Effect of hypertension before beginning dialysis on survival of hemodialysis patients. *American Journal Kidney Disease*. 41:814–821.

Mason, J., Khunti, K., Stone, M., Farooqi, A., & Carr, S. (2008). Educational interventions in kidney disease care: a systematic review of randomized trials. *American Journal of Kidney Diseases* 51, 933–951.

Mazzuchi, N., Carbonell, E., & Fernandez-Cean, J. (2000). Importance of blood pressure control in hemodialysis patient survival. *Kidney International*, 58, 2147-2154.

Ministry of Health, Malaysia. (2004). Renal replacement therapy Clinical Practice Guidelines (2nd ed.).

Molaison, E., & Yadrick, M. (2003). Stages of change and fluid intake in dialysis patients. *Patient Education and Counseling*, 49, 5-12.

Moosa, M.R., Naicker, S., Naiker, I., Pascoe, M., Rensberg, V.B. (2006). Guidelines for the Optimal Care of Patients on Chronic Dialysis in South Africa. South African Renal Society.

Morris, S. B., & DeShon, R. P. (2002). Combining effect size estimates in meta-analysis with repeated measures and independent-groups designs. *Psychological Methods*, 7, 105-125.

Morse, S.A., Dang, Thakur, V., Zhang, R., & Reisin, E. (2003). Hypertension in Chronic Dialysis Patients: Pathophysiology. Monitoring and Treatment. *The American Journal Medical Sciences*, 325(4), 194-201.

Moss, M., & Schwartz, M. (1999). Improving fluid compliance in the hemodialysis population. *Nephrology News Issues*, 13(11), 81-82.

National Kidney Foundation - Kidney Disease Outcome Quality Initiative (NKF-KDOQI) Clinical practice guidelines on Chronic Kidney Disease (2002).

National Kidney Foundation - Kidney Disease Outcome Quality Initiative (NKF-KDOQI) Clinical practice guidelines on Chronic Kidney Disease (2005).

National Kidney Foundation: KDOQI Clinical Practice Guidelines and Clinical Practice Recommendations 2006 (NKF- KDOQI)

National Renal Registry (NRR). (2011). The 19<sup>th</sup> Report of the Malaysian Dialysis and Transplant Registry National Renal Registry (2011), Malaysia.

Newmann, J.M., & Litchfield, W.E. (2005). Adequacy of dialysis: the patient's role and patient concerns. *Seminar Nephrology*, 25, 112-119.

Nicoletta, P., Bernardini, J., & Dacko, C. (2000). Compliance with subcutaneous erythropoietin in peritoneal dialysis patients. *Advanced Peritoneal Dialysis*, 16, 90-92.

Nozaki, C., Oka, M., & Chabooyer, W. (2005). The effects of a cognitive behavioral therapy programme for self-care on hemodialysis patients. *International Journal of Nursing Practice*, 11, 228-236.

Ozkahya,M.,Ok,E.,Toz,H.,Asci,G.,Duman,S.,Basci,A.,Kose,T&Mees,D.E.J.Long term survival rates in haemodialysis treated with strict volume control. *Nephrology Dialysis Transplantation*, 21,3506-3513.

Pang, S.K., Ip, W.Y., & Chang, A. (2001). Psychosocial correlates of fluid compliance among Chinese haemodialysis patients. *Journal of Advanced Nursing*, 35(5), 691-698.

Peixoto,A.J.,&Santos,S.S.F.(2010).Blood pressure management among hemodialysis patients:what had we learn. *Current opinion in Nephrology and Hypertension*, 20, 1561-1566.

Picton, M.L., & Foley, R.N. (2000). Reducing cardiovascular morbidity and mortality from hypertension in end stage disease. *Current Opinion in Nephrology and Hypertension*, 9(5), 497-500.

Pryer,M.D. (2005).The importance of fluid control. *American Association of Kidney Patients – Renalife*, 20 (6), 1-2

Rahman, M., Dixit, A., & Donley, V. (1999). Factors associated with inadequate blood pressure control in hypertensive hemodialysis patients. *American Journal of Kidney Disease*, 33, 498- 506.

Rahman, A., Fu, P., Sehgal, A.R., Smitch, M.C. (2000). Interdialytic weight gain, compliance with dialysis regime and age are independent predictors of blood pressure in hemodialysis patients. *American Journal of Kidney Disease*, 35, 257-265.

Rambod,M.,Peyravi,H., Shokrpour,N.,& Taghi,S.(2010).Dietary and Fluid Adherence in Iranian Hemodialysis Patients. *The Health Care Manager*, 29(4), 359-364.

Richard ,C.J. (2006), Self care management in adults undergoing hemodialysis, *Nephrology Nursing Journal*,33,387-394

Rocco, M.V., Yan, G., Heyka, R.J., Benz, R., & Cheung, A.K. (2001). Risk factor for hypertension in chronic hemodialysis patients: -baseline data from HEMO study. *American Journal of Nephrology*, 21, 280-288.

Saran, R., BraggGresham, J.L., Rayner, H.C., Goodkin, D.A., Keen, M.L., VanDijk, P.C., Korokawa, K., Piera, L., Saito, A.Fukuhara, S., Young, E.W., Hela, P.J., & Port, F.K. (2003). Nonadherence in hemodialysis: Associations with mortality, hospitalization, and practice patterns in the DOPPS. *Kidney International*, 64(1), 254-262.

Sarkar, S.R., Kotanko, P., & Levin, N.W.(2006).Interdialytic Weight Gain: Implications in Hemodialysis Patients. *Seminars in Dialysis*. 19(5), 429-433.

Sathvik, B.S., Mangasuli, S., Narahari, M.G., Gurudev, K.C., & Parthasarathi, G. (2007). Medication knowledge of hemodialysis patients and influence of clinical pharmacist provided education on their knowledge. *Journal of Pharmaceutical Sciences*, 69:232-239.

Schlatter, S., & Ferrans, C.E. (1998).Teaching program effects on high phosphorus levels in patients receiving hemodialysis. *ANNA Journal*, 25, 31-36.

Sehgal, A.R. (2002). Improving Hemodialysis Patient Outcome: A Step-By-Step Approach. *Seminar in Dialysis*, 15(1), 35-37.

Shaldon, S. (2000). Can antihypertensive medications control BP in haemodialysis patients (Letter). *Nephrology Dialysis Transplantation*, 15, 736.

Sun, C.Y., Chang, K.C., Chen, S.H., Chang, C.T., & Wu, M.S. (2008). Patient education: an efficient adjuvant therapy for hyperphosphatemia in hemodialysis patients. *Renal Failure*, 30, 57–62.

Tapolyai,M., Faludi,M., Reti,V., Lengvarszky,Z., Szarvas,T., & Berta,K.(2011). Dialysis Patients' Fluid Overload, Antihypertensive Medications, and Obesity. American Society for Artificial Internal Organs, 57(6), 511-515.

Tsay, S.L., & Healstead, M. (2002). Self-efficacy, depression, and quality of life among patients receiving hemodialysis in Taiwan. *International Journal of Nursing Studies*, 39, 245-251.

Tsay, S.L. (2003). Self –efficacy training for patients with renal - stage renal disease. *Journal of Advanced Nursing*, 43(4), 379-375.

Thomas, N. (2002). *Renal Nursing* (2<sup>nd</sup> ed.). London: Bailliere Tindall.

Tomson, C.R.V. (2001). Advising dialysis patients to restrict fluid intake without restricting sodium intake is not based on evidence and is a waste of time. *Nephrology Dialysis Transplantation*, 16, 1538-1542.

Valdez, R. (2003). Dealing with noncompliant dialysis patients: 10 steps to achieve compliance. *Nephrology News & Issues*, 4, 65-69.

Van Camp,Y.P., Huybrechts,S.A., Rompaey,B.A., & Elseviers,M.M. (2011). Nurse-led education and counselling to enhance adherence to phosphate binders. *Journal of Clinical Nursing*, 21, 1304–1313.

Vernaglione, L., Nosell, V., Chimienti, S.(2010). Correlates of Blood Pressure and Predictors of Cardiovascular Mortality in Haemodialysis Patients: A Clinical Experience High Blood Pressure & Cardiovascular Prevention. 17(2):65-72.

Vlaminck, H., Maes, B., Jacobs, A., Reyntjens, S., & Evers, G. (2001). The dialysis diet and fluid non-adherence questionnaire: validity testing of a self-report instrument for clinical practice. *Journal of Clinical Nursing*, 10(5), 707-714.

Welch,J.K., & Thomas-Hawkins. C, (2005). Psycho-education strategies to promote fluid adherence in adult hemodialysis patients: a review of international studies . *International Journal of Nursing Studies*, 42, 597-608

Welch, J.L., Perkins, S.M., Johnson, C.S., & Kraus, M.A. (2006). Patterns of interdialytic weight gain during the first year of hemodialysis. *Nephrology Nursing Journal* 33, 493-499.

Weir, M.R. (2010). Hypervolemia and Blood Pressure Powerful Indicators of Increased Mortality Among Hemodialysis Patients. *Hypertension* , 56, 341-345.

Yoon,S.J. ,Conway,J.,McMillan.(2006). An exploration of the concept of patient education: Implications for the development of educational programmes for relapsed post bone- marrow transplantations patients and their families in Korea. *International Journal of Nursing Practice*, 12(3), 129-135.

Zager, G., Nikolic, J., Brown, R.H., Campbell, M.A., Hunt, W.C., Peterson, D., Van, S.J Levey, A., Meyer, K.B., Klag, M.J., Johnson, H.K., Clark, E., Sadler, J.H., & Teredesai,P. (1998). "U"curve association of blood pressure and mortality in hemodialysis patients. *Kidney International*, 54, 561-569.

Zerbi, S. P. & Luciano, A. (2010). Hypertension in Hemodialysis Patients.*Current Hypertension Reviews*. 6(1):44-54.

Zrinyi,M.,Juhasz,M.,Balla,J.,Katona,E.,Ben,T.,Kakuk,G.Pall,D.(2003). Dietary self - efficacy: determinant of compliance behavior and biochemical outcomes in haemodialysis patients. *Nephrology Dialysis Transplantation*, 18(9), 1869-1873.