

## LIST OF FIGURES

Figure 2.1	Parts of the mushroom	16
Figure 2.2	Some edible mushrooms that can be found in Malaysia	20
Figure 2.3	Commonly available edible <i>Pleurotus</i> spp. mushroom locally grown in Malaysia.	25
Figure 2.4	Commercially available oyster mushroom chips in Malaysia	28
Figure 3.1	Extraction and fractionation of fresh fruiting bodies <i>P. sajor-caju</i> .	39
Figure 3.2	A column chromatography setup	42
Figure 3.3	Dialysis of polysaccharide extract (GE) using a DEAE cellulose dialysing tube against distilled water for 4 hr	43
Figure 3.4	A photograph showing the fruiting bodies of <i>P. sajor-caju</i>	45
Figure 3.5	Yield of <i>P. sajor-caju</i> extracts after extraction and fractionation from fresh fruiting bodies	46
Figure 3.6	Mass spectrum of 2,4-ditertbutylphenol	49
Figure 3.7	Mass spectrum for ethyl myristate	50
Figure 3.8	Mass spectrum of methyl palmitate	50
Figure 3.9	Mass spectrum of ethyl palmitate	51
Figure 3.10	Mass spectrum of methyl linoleate	51
Figure 3.11	Mass spectrum of methyl stearate	52
Figure 3.12	Mass spectrum of ethyl linoleate	52
Figure 3.13	Mass spectrum of ethyl stearate	53
Figure 3.14	McLafferty rearrangement of methyl palmitate	54
Figure 3.15	Mass spectrum of palmitic acid	56
Figure 3.16	Mass spectrum of oleic acid	56
Figure 3.17	Mass spectrum of ethyl oleate	57
Figure 3.18	Mass spectrum of ethyl linolenate	57
Figure 3.19	Mass spectrum of ergosta-5,8,22-trien-3-ol	58
Figure 3.20	Mass spectrum of 5,6-dihydroergosterol	59
Figure 3.21	Mass spectrum of neoergosterol	59
Figure 3.22	Mass spectrum of ergosta- 5,8-dien-3-ol	60
Figure 3.23	Mass spectrum of ergosta-7- en-3-ol	60

Figure 3.24	Mass spectrum of myristic acid	61
Figure 3.25	Mass spectrum of pentadecanoic acid	62
Figure 3.26	Mass spectrum of linoleic acid	62
Figure 3.27	Mass spectrum of ergosta- 5,8(14)-dien-3-ol	63
Figure 3.28	Mass spectrum of ergost- 5-en-3-ol	63
Figure 3.29	Structure of ergosterol identified using GC-MS and NMR analysis	65
Figure 3.30	Mass spectrum of 4-hydroxybenzaldehyde	66
Figure 3.31	Mass spectrum of benzeneacetamide	67
Figure 3.32	Mass spectrum of nicotinamide	67
Figure 3.33	Mass spectrum of cinnamic acid	68
Figure 3.34	Yield of crude polysaccharides of <i>P. sajor-caju</i> after isolation and purification from hot water extraction	68
Figure 4.1	Correlation between antioxidant assays and total phenolic content; (a) total phenolic content and ferric reducing antioxidant power assay and (b) total phenolic content and trolox equivalents antioxidant capacity assay.	98
Figure 5.1	The series of pictures shows (a) mature adipocytes and (b) mature adipocytes stained with Oil Red O dye.	117
Figure 5.2	TaqMan® gene expression assay chemistry	125
Figure 5.3	Real Time RT-PCR instrument by Applied Biosystem with StepOnePlus® software	126
Figure 5.4	The percentages of proliferation of preadipocytes cells after treatment with <i>P. sajor-caju</i> extracts (a) after 24 hours treatment and (b) after 48 hours treatment	129
Figure 5.5	Lipogenic effect of various concentrations of <i>P. sajor-caju</i> extracts on adipocytes measured using Oil Red O quantification.	131
Figure 5.6	Lipolytic effect of various concentrations of <i>P. sajor-caju</i> extracts on 3T3-L1 adipocytes. (a) quantification of glycerol (b) quantification of free fatty acids	133
Figure 5.7	Effects of GE and insulin on gene expression in 3T3-L1 adipocytes.	137
Figure 5.8	Regulation of lipolysis in adipocytes and possible involvement of GE.	140
Figure 6.1	Instruments used to study oral glucose tolerance test (OGTT).	150

Figure 6.2	The experimental design and analytical methods for the <i>in vivo</i> study	152
Figure 6.3	Effects of GE on the changes of body weight and lipid profile of C57BL/6J mice fed on a high-fat diet or normal diet	159
Figure 6.4	The size of mouse from each group after 15 weeks of experiment	160
Figure 6.5	Effects of GE on oral glucose tolerance test (OGTT) of C57BL/6J mice fed on a high-fat diet or normal diet	164
Figure 6.6	Effects of GE on fasting plasma insulin, fasting blood glucose levels and HOMA-IR value of C57BL/6J mice fed on a high-fat diet or normal diet	167
Figure 6.7	Effects of GE and metformin on protein carbonyl content (AOPP), lipid hydroperoxides, uric acid and DNA damage levels in the urine samples of C57BL/6J mice fed on a high-fat diet or normal diet	170
Figure 6.8	Inflammation and altered secretion of inflammatory markers in HFD group	186
Figure 7.1	Overview of important findings and implication from the present research	195