

## REFERENCES

- Agarwal, R. & Chase, S.D. (2002). Rapid, flurometric-liquid chromatographic determination of malondialdehyde in biological samples. *Journal of Chromatography*, 775, 121–126.
- Agrawal, R. P., Chopra, A., Lavekar, G.S., Padhi, M.M., Srikant, M., Ota, S. & et al. (2010). Effect of oyster mushroom on glycemia, lipid profile and quality of life in type 2 diabetic patients. *Australian Journal of Medical Herbalism*, 22, 50–54.
- Ahmadian, M., Wang, Y. & Sul, H.S. (2010). Lipolysis in adipocytes. *The International Journal of Biochemistry & Cell Biology*, 42, 555–559.
- Akbar, D.H. (2003). Effect of metformin and sulfonylurea on C-reactive protein level in well-controlled type 2 diabetics with metabolic syndrome. *Endocrine*, 20, 215–218.
- Akehi, Y., Anzai, K., Katsuta, H., Yoshida, R., Ohkubo, K., Yamashita, T. & et al. (2008). Adverse effects of obesity on  $\beta$ -cell function in Japanese subjects with normal glucose tolerance. *Obesity Research & Clinical Practice*, 2, 195–202.
- Amarowicz, R., Pegg, R.B., Moghaddam, P.R., Barl, B. & Weil, J.A. (2004). Free-radical scavenging capacity and antioxidant activity of selected plant species from the Canadian Prairies. *Food Chemistry*, 84(4), 551–562.
- Ando, K. & Fujita, T. (2009). Metabolic syndrome and oxidative stress. *Free Radical Biology & Medicine*, 47, 213–218.
- Aliaga, M.J.M., Stanhope, K.L., Gregoire, F.M., Warden, C.H. & Havel, P.J. (2003). Effects of inhibiting transcription and protein synthesis on basal and insulin stimulated leptin gene expression and leptin secretion in cultured rat adipocytes. *Biochemical & Biophysical Research Communications*, 307, 907–914.
- Antras, J., Lasnier, F. & Pairault, J. (1991). Beta-adrenergic-cyclic-AMP signaling pathway modulate cell function at the transcriptional level in 3T3-F442A adipocytes. *Molecular & Cellular Endocrinology*, 82, 183–190.
- Arner, P. (2003). The adipocyte in insulin resistance: key molecules and the impact of thiozolidinediones. *Trends in Endocrinology & Metabolism*, 14, 137–145.

- Astrup, A. & Finer, N. (2000). Redefining type 2 diabetes: 'diabesity' or obesity dependent diabetes mellitus. *Obesity Review*, 1, 57–59.
- Ayerdi, S.G.S., Brenes, A. & Goni, I. (2009). Effect of grape antioxidant dietary fiber on the lipid oxidation of raw and cooked chicken hamburgers. *LWT-Food Science & Technology*, 42, 971–976.
- Bahl, N. (1994). Handbook on Mushrooms. Oxford & IBH Publishing Co. Ltd. New Delhi.
- Bai, L., Pang, W.J., Yang, Y.J. & Yang, G.S. (2008). Modulation of Sirt1 by resveratrol and nicotinamide alters proliferation and differentiation of pig preadipocytes. *Molecular & Cell Biochemistry*, 307, 129–140.
- Bailey, C.J. (1999). Insulin resistance and antidiabetic drugs. *Biochemical Pharmacology*, 58, 1511–1520.
- Bakker, W., Eringa, E.C., Sipkema, P. & Hinsberg, V.W.M. (2009). Endothelial dysfunction and diabetes: roles of hyperglycemia, impaired insulin signaling and obesity. *Cell Tissue Research*, 335, 165–189.
- Balagopal, P., Graham, T.E., Kahn, B.B., Altomare, A., Funanage, V. & George, D. (2007). Reduction of elevated serum retinol-binding protein 4 in obese children lifestyle intervention: association with sub-clinical inflammation. *Journal of Clinical Endocrinology & Metabolism*, 7, 2006–2012.
- Bano, Z. & Rajaratnam, S. (1988). Pleurotus mushrooms. Chemical composition, nutritinal Value, Post-Harvest Physiology, Preservation and Role as Human Food. *CRC Critical Reviews in Food Science & Nutrition*, 27(2), 87–158.
- Barnea, M., Shamay, A., Stark, A.H. & Madar, Z. (2006). A high-fat diet has a tissue-specific effect on adiponectin and related enzyme expression. *Obesity*, 14, 2145–2153.
- Barros, L., Duenas, M., Ferreira, I.C.F.R., Baptista, P., Buelga, C.S. (2009). Phenolic acids determination bu HPLC-DAD-ESI/MS in sixteen different Portuguese wild mushrooms species. *Food Chemistry & Toxicology*, 47, 1076–1079.

- Barros, L., Baptista, P., Correia, D.M., Casal, S., Oliveira, B. & et al. (2007). Fatty acid and sugar compositions and nutritional value of five wild edible mushrooms from Northeast Portugal. *Food Chemistry*, 105, 140–145.
- Benzie, I.F.F. & Strain, J.J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of ‘antioxidant power’: The FRAP assay. *Analytical Biochemistry*, 74, 2268–2277.
- Benzie, I.F.F. & Strain, J.J. (1999). Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods in Enzymology*, 299, 15–27.
- Bhattaram, V.A., Graefe, U., Kohlert, C., Veit, M. & Derendorf, H. (2002). Pharmacokinetics and bioavailability of herbal medicinal products. *Phytomedicine*, 9, 1–6.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72, 248–54.
- Broncano, J.M., Petron, M.J., Parra, V. & Timon, M.L. (2009). Effects of different cooking methods on lipid oxidation and formation of free cholesterol oxidation products (COPs) in *Latissimus dorsi* muscle of Iberian pigs. *Meat Science*, 83, 431–437.
- Brun, R.P., Kim, J.B., Hu, E. & Spiegelman, B.M. (1997). Peroxisome proliferator-activated receptor gamma and the control of adipogenesis. *Current Opinion on Lipidol*, 8, 212–218.
- Bruun, J.M., Pedersen, S.B., Kristensen, K. & Richelsen, B. (2002). Effects of pro-inflammatory cytokines and chemokines on leptin production in human adipose tissue in vitro. *Molecular & Cellular Endocrinology*, 190, 91–99.
- Bountagkidou, O.G., Ordoudi, S.A. & Tsimidou, M.Z. (2010). Structure-antioxidant activity relationship study of natural hydroxybenzaldehydes using *in vitro* assays. *Food Research International*, 43, 2014–2019.
- Boer, V.C., Dihal, A.A., Woude, H., Art, I.C., Wolffram, S., Alink, G.M. & et al. (2005). Tissue distribution of quercetin in rats and pigs. *Journal of Nutrition*, 135, 1617–1618.

- Bonnard, C., Durand, A., Vidal, H. & Rieusset, J. (2008). Changes in adiponectin, its receptor and AMPK activity in tissues of diet induced-diabetic mice. *Diabetes & Metabolism*, 34, 52–61.
- Borchers, A.T., Keen, C.L. & Gershwin, M.E. (2004). Mushrooms, Tumors and Immunity: An Update. *Experimental Biology*, 229, 393–406
- Caglarirmak, N. (2007). The nutrients of exotic mushrooms (*Lentinus edodes* and *Pleurotus* species) and an estimated approach to the volatile compounds. *Food Chemistry*, 105, 1188–1194.
- Caillet, S., Yu, H., Lessard, S., Lamoureux, G., Ajdukovic, D., & Lacroix, M. (2007). Fenton reaction applied for screening natural antioxidants. *Food Chemistry*, 100(2), 542–552.
- Castrillejo, V.M., Romero, M.M., Esteve, M., Ardevol, A., Blay, M., Blade, C. & et al. (2011). Antioxidant effect of grape seed procyanidin extract and oleoyl-estrone in obese Zucker rats. *Nutrition*, doi:10.1016/j.nut.2010.12.010.
- Chang, S.T., Chen, P.F. & Chang, S.C. (2001). Antibacterial activity of leaf essential oils and their constituents from *Cinnamomum osmophloeum*. *Journal of Ethnopharmacology*, 77, 123–127.
- Chang, S.T. (1999). Global impact of edible and medicinal mushrooms on human welfare in the 21<sup>st</sup> century: Non-green revolution. *International Journal of Medicinal Mushrooms*, 1, 1–8.
- Chang, S.T. & Bushwell, J.A. (1999). *Ganoderma lucidum* (Curt.; Fr) P. Karst (Aphyllphoromycetidae)- a mushrooming medicinal mushroom. *International Journal of Medicinal Mushrooms*, 1, 139–146.
- Chang, S.T. (1993). Mushrooms biology: The impact on mushrooms production and mushrooms products (Ed.). *Mushroom biology and mushrooms products*, Hong Kong: Chinese University Press, 3–20.
- Chaves, V.E., Frasson, D. & Kawashita, N.H. (2011). Several agents and pathways regulate lipolysis in adipocytes. *Biochimie*, 93, 1631–1640.

- Cheung, L.M., Cheung, P.C.K. & Ooi, V.E.C. (2003). Antioxidant activity and total phenolics of edible mushrooms extracts. *Food Chemistry*, 81, 249–255.
- Cheung, L.M & Cheung, P.C.K. (2005). Mushrooms extracts with antioxidant activity against lipid peroxidation. *Food Chemistry*, 89, 403–409.
- Chirinang, P. & Intarapichet, K.O. (2009). Amino acids and antioxidant properties of the oyster mushrooms, *Pleurotus ostreatus* and *Pleurotus sajor-caju*. *ScienceAsia*, 35, 326–331.
- Cho, S., Choi, Y., Park, S. & Park, T. (2011). Carvacrol prevent diet induce obesity by modulating gene expressions involved in adipogenesis and inflammation in mice fed with high - fat diet. *Journal of Nutritional Biochemistry*, doi: 10. 1016/j.jnutbio. 2010.11.016.
- Choi, H., Eo, H., Park, K., Jin, M., Park, E.J., Kim, S.H. & et al. (2007). A water soluble extract from *Curcubita mocshata* shows anti-obesity effects by controlling lipid metabolism in high fat diet-induced obesity mouse model. *Biochemical & Biophysical Research Communications*, 359, 419–425.
- Choi, Y., Lee, S.M., Chun, J., Lee, H.B. & Lee, J. (2006). Influence of heat treatment on antioxidant activities and polyphenolic compounds of Shiitake (*Lentinus edodes*) mushroom, *Food Chemistry*, 99, 381–387.
- Chu, K.T., Xia, L. & Ng, T.B. (2005). Pleurotin, an antifungal peptide from oyster mushroom. *Peptides*, 26, 2098–2103.
- Cimerman, N.G. (1999). Medicinal value of the genus *Pleurotus* (Fr.) P.Karst. (Agaricales s.i., Basidiomycetes). *International Journal of Medicinal Mushrooms*, 1, 69–80.
- Clapham, J.C., Arch, J.R.S. & Tadayyon, M. (2001). Anti-obesity drugs: a critical review of current therapies and future opportunities. *Pharmacology & Therapeutics*, 89, 81–121.
- Clarke, S.D., Turini, M. & Jump, D. (1997). Polyunsaturated fatty acids regulate lipogenic and peroxisomal gene expression by independent mechanisms. *Prostaglandins, Leukotrienes & Essential Fatty Acids*, 57(1), 65–69.

- Coate, K.C. & Huggins, K.W. (2010). Consumption of high glycemic index diet increases abdominal adiposity but does not influence adipose tissue pro-oxidant and antioxidant gene expression in C57L/6J mice. *Nutrition Research*, 30, 141–150.
- Coudray, C.F., Sutra, T., Fouret, G., Ramos, J., Cabello, C.W., Cabello, G. & et al. (2009). Oxidative stress in rats fed a high fat high sucrose diet and preventive effect of polyphenols: involvement in mitochondrial and NAD(P)H oxidase systems. *Free Radical Biology & Medicine*, 46, 624–632.
- Courturier, C., Janvier, B., Girlich, D., Bereziat, G. & Mangeney, M.A. (1998). Effects of caffeine on lipoprotein gene expression during the adipocyte differentiation process, *Lipids*, 33, 455–460.
- Crisan, E.V. & Sands, A. (1978). Nutritional Value. In: *The Biology and Cultivation of Edible Mushrooms*. Academic Press, 137–165.
- Crevel, R.W., Friend, J.V., Goodwin, B.F. & Parish, W.E. (1992). High fat diets and the immune response of C57Bl mice. *British Journal of Nutrition*, 67, 17–26.
- Crowwell, W.L. & Otvos, J.D. (2004). Low density lipoprotein particles number and risk for cardiovascular disease. *Current Artherosclerosis. Research*, 6, 381–387.
- Cvetkovic, T., Mitic, B., Lazarevic, G., Vlahovic, P., Antic, S. & Stefanovic, V. (2009). Oxidative stress parameters as possible urine markers in patients with diabetic nephropathy. *Journal of Diabetes & its Complications*, 23, 337–342.
- Daker, M., Abdullah, N., Vikineswary, S., Goh, P.C. & Kuppusamy, U.R. (2008). Antioxidant from maize and maize fermented by *Marasmiellus* sp. as stabilizer of lipid rich foods. *Food Chemistry*, 107, 1092–1098.
- Daizo, A., Egashira, Y. & Sanada, H. (2005). Suppressive effect of corn bran hemicellulose on liver injury induced by D-galactosamine in rats. *Nutrition*, 21, 1044–1051.
- Dallas, C., Gerbi, A., Tenca, C., Juchaux, F. & Bernard, F.X. (2008). Lipolytic effects of a polyphenolic citrus dry extract of red orange, grapefruit, orange (SINETROL) in human body fat adipocytes. Mechanism of action by inhibition of cAMP – phosphodiesterase (PDE). *Phytomedicine*, 15, 783–792.

- Dandona, P., Aljada, A. & Bandyopadhyay, A. (2004). Inflammation: the link between insulin resistance, obesity and diabetes. *Trends Immunology*, 25, 4–7.
- Despres, J.P. (2003). Potential contribution of metformin to the management of cardiovascular disease risk in patients with abdominal obesity, the metaolic syndrome and type 2 diabetes. *Diabetes Metabolism*, 29, 6S54–6S61.
- Dey, B., Bhunia, S.K., Maity, K.K., Patra, S., Mandal, S., Maiti, S. & et al. (2010). Chemical analysis of an immunoenhancing water-soluble polysaccharides of an edible mushroom, *Pleurotus florida* blue variant. *Carbohydrate Research*, 345, 2736–2741
- Dizdaroglu, M., Jaruga, P., Birincioglu, M. & Rodriguez, H. (2002). Free radical induced damage to DNA: mechanisms and measurement. *Free Radical Biology & Medicine*, 32, 1102–1115.
- Donahoo, W.T., Stob, N.R., Ammon, S., Levin, N. & Eckel, R.H. (2011). Leptin increases skeletal muscle lipoprotein lipase and postprandial lipid metabolism in mice. *Metabolism Clinical & Experimental*, 60, 438–443.
- Dore, C.M.P.G., Azevedo, T.C.G., Souza, M.C.R., Rego, L.A., Dantas, J.C.M., Silva, F.R.F. & et al. (2007). Antiinflammatory, antioxidant and cytotoxic action of  $\beta$ -glucan rich extract from *Geastrum saccatum* mushroom. *International Immunopharmacology*, 7, 1160–1169.
- Du, G., Li, M., Ma, F. & Liang, D. (2009). Antioxidant capacity and the relationship with polyphenol and vitamin C in *Actidinia* fruits. *Food Chemistry*, 113, 557–562.
- Elmastas, M., Isildak, O., Turkecul, I. & Temur, N. (2007). Determination of antioxidant activity and antioxidant compounds in wild edible mushrooms, *Journal of Food Composition & Analysis*, 20, 337–345.
- El-Moselhy, M.A., Taye, A., Sharkawi, S.S. & El-Sisi, S.F.I. (2011). The antihyperglycemic effect of curcumin in high-fat diet fed rats. Role of TNF- $\alpha$  and free fatty acids. *Food & Chemical Toxicology*, 49(5), 1129–1140.
- Enman, J., Rova, U., & Berglund, K. A. (2007). Quantification of the bioactive compound eritadenine in selected strains of shiitake mushroom (*Lentinus edodes*). *Journal of Agricultural & Food Chemistry*, 55, 1177–1180.

- Esterbauer, H. & Cheeseman, K.H. (1990). Determination of aldehydic lipid peroxidation products: malonaldehyde and 4-hydroxynonenal. *Methods Enzymology*, 186, 407–421.
- Fardet, A., Canlet, C., Gottardi, G., Lyan, B., Llorach, R., Ramsey, C & et al. (2007). Supplementation of diets with the black rice pigment fraction attenuates atherosclerotic plaque formation in apolipoprotein E deficient mice. *Journal of Nutrition*, 133, 741–751.
- Fasshauer, M., Klein, J., Neumann, S., Eszlinger, M. & Paschke, R. (2002). Hormonal regulation of adiponectin gene expression in 3T3-L1 adipocytes. *Biochemical and Biophysical Research Communications*, 290, 1084–1089.
- Fedotov, O.V. (2001). Antioxidizing activity of mycelium of mushroom stocks *Pleurotus* (Fr.) Kumm. and *Flammulina* (Curt.:Fr.) Sing. *International Journal of Medicinal Mushrooms*, 3(2-3), 1.
- Ferrante, A.W. (2007). Obesity induced inflammation: a metabolic dialogue in the language of inflammation. *Journal of International Medicine*, 262, 408–414.
- Fielding, B.A. & Frayn, K.N. (1998). Lipoprotein lipase and the disposition of dietary fatty acids. *The British Journal of Nutrition* 80, 495–502.
- Frankel, E.N. & Meyer, A.S. (2000). The problems of using one-dimensional methods to evaluate multifunctional food and biological antioxidants. *Journal of Science Food Agricultural*, 80, 1925–1941.
- Fruebis, J., Tsao, T.S., Javorschi, S., Ebbets-Reed, D., Erickson, M.R. & Yen, F.T. (2001). Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natural Academia Science*, 98, 2005–2010.
- Frode, T.S. & Medeiros, Y.S. (2008). Animal model to test drugs with potential antidiabetic activity. *Journal of Ethnopharmacology* 115, 173–183.
- Fong, J.C. (1990). The effect of chronic fatty acid treatment on lipolysis in 3T3-L1 adipocytes. *Biochemical & Biophysical Research Communications*, 171(1), 46–52.



- Fu, L., Isobe, K., Zen, Q., Suzukawa, K., Takekoshi, K. & Kawakami, Y. (2007).  $\beta$ -adrenoceptor agonists downregulate adiponectin, but upregulate adiponectin receptor 2 and tumor necrosis factor  $\alpha$  expression in adipocytes. *European Journal of Pharmacology*, 569, 155–162.
- Furukawa, S., Fujita, T., Shimabukuro, M., Iwaki, M., Yamada, Y., Nakajima, Y. & et al. (2004). Increased oxidative stress in obesity and its impact on metabolic syndrome. *Journal of Clinical Investigation*, 114, 1752–1761.
- Fukushima, M., Ohashi, T., Fujiwara, Y., Sonoyama, K. & Nakano, M. (2001). Cholesterol lowering effects of Maitake fiber, Shiitake fiber and Enokitake fiber in rats. *Experimental Biology & Medicine*, 8, 758–765.
- Galli, F., Piroddi, M., Annetti, C., Aisa, C., Floridi, E. & Floridi, A. (2005). Oxidative stress and reactive oxygen species. *Contribution Nephrology*, 149, 240–260.
- Gao, J.M., Wang, M., Liu, L.P., Wei, G.H., Zhang, A.L., Draghici, C. & Konishi, Y. (2007). Ergosterol peroxides as phospholipase A<sub>2</sub> inhibitors from the fungus *Lactarius hatsudake*. *Phytomedicine*, 14, 821–824.
- Gern, R.M.M., Wisbeck, E., Rampinelli, J.R., Ninow, J.L. & Furlan, S.A. (2008). Alternative medium for production of *Pleurotus ostreatus* biomass and potential antitumor polysaccharides. *Bioresource Technology*, 99, 76–82.
- Ghoreishi, S.M., Mardani, E. & Ghaziaskar, H.S. (2011). Separation of  $\gamma$ -linolenic and other polyunsaturated fatty acids from *Boraginaceae* via supercritical CO<sub>2</sub>. *Journal of Separation Science*, 34, 233–240.
- Glueck, C.H., Aregawi, D., Agloria, M., Winiarska, M., Sieve, L. & Wang, P. (2006). Sustainability of 8% weight loss, reduction of insulin resistance, and amelioration of atherogenic-metabolic risk factors over 4 years by metformin-diet in women with polycystic ovary syndrome. *Metabolism Clinical & Experimental*, 55, 1582–1589.
- Griffin, S. & Bhagooli, R. (2004). Measuring antioxidant potential in corals using the FRAP assay. *Journal of Experimental Marine Biology & Ecology*, 302, 373–414.
- Green, H., & Kehinde, O. (1974). Sublines of mouse 3T3 cells that accumulate lipid. *Cell*, 1, 113–116.

- Gomes, T., Delcuratolo, D.,Paradiso, V.M., Summo, C. & Caponio, F. (2011). Pro-oxidant activity of oxidized triacylglycerols in olive oil and comparison with pro-oxidant action of polar triacylglycerol oligopolymers. *LWT-Food Science & Technology*, 44, 1236–1239.
- Graham, T.E., Yang, Q., Bluher, M., Hammarstedt, A., Ciaraldi, T.P., Henry, R.R. & et al. (2006). Retinol-binding protein 4 and insulin resistance in lean, obese and diabetic subjects. *England Journal of Medicine*, 354, 2552-2563.
- Granneman, J.G., Moore, H.P., Granneman, R.L., Greenberg, A.S., Obin, M.S. & Zhu, Z. (2007). Analysis of lipolytic protein trafficking and interactions in adipocytes. *Journal Biological Chemistry*, 282, 5726–5735.
- Gravina, H.D., Tafuri, N.F., Junior, S.A., Fietto, J.L.R., Oliveira, T.T., Diaz, M.A.N. & Almeida, M.R. (2010). *In vitro* assessment of antiviral potential of trans-cinammic acid, quercetin and morin against *equid herpesvirus 1*. *Research in Veterinary Science*, 91, 158–162
- Gokalp, D., Bahceci, M.,Ozmen, S., Arikan, S., Tuzcu, A., & Danis, R. (2008). Adipocyte volume and levels of adipokines in diabetes and obesity. *Diabetes Metabolism Syndrome;Clinical Research & Review*, 2, 253–258.
- Golay, A. & Ybarra, J. (2005). The link between obesity and and type 2 diabetes. *Best Practice & Research Clinical Endocrinology & Metabolism*, 19, 649–663.
- Gunde-Cimerman, N. & Cimerman, A. (1995). *Pleurotus* fruiting bodies contain the inhibitor of 3-hydroxy-3-methylglutaryl-coenzymeA reductase-lovastatin. *Experimental Mycology*, 19, 1–6.
- Guo, W., Wong, S., Xie, W., Lei, T. & Luo, Z. (2007). Palmitate modulates intracellular signaling, induces endoplasmic reticulum stress, and causes apoptosis in mouse 3T3-L1 and rat primary preadipocytes. *American Journal of Physiology*, 293(2), 576–586.
- Guo, C., Yang, J., Wei, J., Li, Y., Xu, J. & Jiang, Y. (2003). Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutrition Research*, 23, 1719–1726.
- Guo, X. & Liao, K. (2000). Analysis of gene expression profiling during 3T3-L1 preadipocyte differentiation. *Gene*, 251, 45–53.

- Guzman, G. (2000). Genus *Pleurotus* (Jacq.:Fr.) P.Kumm. (Agraricomycetideae): Diversity, taxonomy problems and cultural and traditional medicinal uses. *International Journal of Medicinal Mushrooms*, 2, 95-123.
- Hajer, G.R., Haeften, T.W. & Visseren, F.L.J. (2008). Adipose tissue dysfunction in obesity, diabetes, and vascular diseases. *European Heart Journal*, 29(4), 2959–2971.
- Hamza, N., Berke, B., Cheze, C., Agli, A.N., Robinson, P., Gin, H. & et al. (2010). Prevention of type 2 diabetes induced by high fat diet in the C57BL/6j mouse by two medicinal plants used in traditional treatment of diabetes in the east of Algeria. *Journal of Ethnopharmacology*, 128, 513–518.
- Han, C., Yuan, J., Wang, Y. & Li, L. (2006). Hypoglycemic activity of fermented mushroom of *Coprinus comatus* rich in vanadium. *Journal of Trace Elements in Medicine & Biology*, 20, 191–196.
- Harborne, J.B. (1973). *Phytochemical methods: A guide to modern techniques of plant analysis*. Publisher: Chapman and Hall, London.
- Hardie, L.J., Guilhot, N. & Trayhurn, P. (1996). Regulation of leptin production in cultured mature white adipocytes. *Hormone Metabolite Research*, 28, 685–689.
- Hasani, H.A. & Joost, H.G. (2005). Nutrition diet induced changes in gene expression in white adipose tissue. *Clinical Endocrinology & Metabolism*, 19, 589–603.
- Haslam, D. (2010). Obesity and diabetes: the links and common approaches. *Primary Care Diabetes*, 4, 105–112.
- Harwood, H.J. (2012). The adipocyte as an endocrine organ in the regulation of metabolic homeostasis. *Neuropharmacology*, doi:10.1016/j.neuropharm.2011.12.010
- Hays, N.P., Galesseti, P.R. & Coker, R.H. (2008). Prevention and treatment of type 2 diabetes: current role of lifestyle, natural product, and pharmacological interventions. *Pharmacology & Therapeutics*, 118, 181-191.
- Heleno, S.A., Barros, L., Sousa, M.J., Martins, A. & Ferreira, I.C.F.R. (2010). Tocopherols composition of Portuguese wild mushrooms with antioxidant capacity. *Food Chemistry*, 119, 1443–1450.

- Higuchi, R., Fockler, C., Dollinger, G., & Watson, R. (1993). Kinetic PCR analysis: real time monitoring of DNA amplification reactions. *Biotechnology*, 11(9), 1026-1030.
- Hu, J.H., Lee, D.U., Lee, J.T., Kim, J.S., Yong, C.S., Kim, J.A., Ha, J.S. & Huh, K. (2001). 4-hydroxybenzaldehyde from *Gastrodia elata* B1. is active in the antioxidant and GABAergic neuromodulation of the rat brain. *Journal of Ethnopharmacology*, 73, 329–333.
- Hong, J.H. & Lee, I.N. (2009). Effects of Artemisia ethyl acetate fraction on oxidative stress and antioxidant enzyme in high fat diet induced obese mice. *Chemico-Biological Interactions*, 179, 88–93.
- Hsu, H.K., Yang, Y.C., Hwang, J.H. & Hong, S.J. (2003). Effects of Toona sinensis leaf extract on lipolysis in differentiated 3T3-L1 adipocytes. *Koahsiung Journal of Medical Sciences*, 19, 385–390.
- Hsu, C.L. & Yen, G.C. (2007). Effects of flavanoids and phenolic acids on the inhibition of adipogenesis in 3T3-L1 adipocytes. *Journal of Agricultural & Food Chemistry*, 55, 8404–8410.
- Hu, S.H., Liang, Z.C., Chia, Y.C., Lien, J.L., Chen, K.S., Lee, M.Y. & Wang, J.C. (2006). Antihyperlipidemic & antioxidant effects of extracts from *Pleurotus citrinopileatus*. *Journal of Agricultural & Food Chemistry*, 54, 2103-2110.
- Huang, D., Ou, B., & Prior, R.L. (2005). The chemistry behind antioxidant capacity assays. *Journal of Agricultural & Food Chemistry*, 53, 1841–1856.
- Huang, Y.W., Liu, Y., Dushenkov, S., Ho, C.T. & Huang, M.T. (2009). Anti-obesity effects of epigallocatechin-3-gallate, orange peel extract, black tea extract, caffeine and their combinations in a mouse model. *Journal of Functional Foods*, 1, 304–310.
- Huang, Y.C., Chang, W.L., Huang, S.F., Lin, C.Y., Lin, H.C. & Chang, H.C. (2010). Pachymic acid stimulates glucose uptake through enhanced GLUT4 expression and translocation. *European Journal of Pharmacology*, 648, 39–49.
- Hunt, E. Plurimutillin antibiotics. *Drugs for Future*, 25, 1163–1170.

- Huypens, P., Quartier, E., Pipeleers, D. & Castele, M.V. (2005). Metformin reduces adiponectin protein expression and release in 3T3-L1 adipocytes involving activation of AMP activated protein kinase, *European Journal of Pharmacology*, 518, 50–59.
- Hwang, S.L., Kim, H.N., Jung, H.H., Kim, J.E., Choi, D.K., Hur, J.M. & et al. (2008). Beneficial effects of beta-sitosterol on glucose and lipid metabolism in L6 myotube cells are mediated by AMP-activated protein kinase. *Biochemical & Biophysical Research Communications*, 377, 1253–1258.
- Iwalokun, B.A., Usen, U.A., Otunba, A.A. & Olukoya, D.K. (2007). Comparative phytochemical evaluation, antimicrobial and antioxidant properties of *Pleurotus ostreatus*. *African Journal of Biotechnology*, 6, 1732–1739.
- Jasinghe, V.J., Perera, C.O. & Sablani, S.S. (2007). Kinetics of the conversion of ergosterol in edible mushrooms. *Journal of Food Engineering*, 79, 864–869.
- Jayakumar, T., Thomas, P.A. & Geraldine, P. (2007). Protective effect of an extract of the oyster mushroom, *Pleurotus ostreatus*, on antioxidant of major organs of aged rats. *Experimental Gerontology*, 42, 183–191.
- Jayakumar, T., Thomas, P.A., Sheu, J.R. & Geraldine, P. (2011). *In-vitro* and *in-vivo* antioxidant effects of the oyster mushroom *Pleurotus ostreatus*. *Food Research International*, 44, 851–861.
- Jayaprakasha, G.K., Singh, R.P. & Sakariah, K.K. (2001). Antioxidant activity of grape seed (*Vitis vinifera*) extracts on peroxidation models *in vitro*. *Food Chemistry*, 73, 285–290.
- Jeong, S.C., Jeong, Y.T., Yang, B.K., Islam, R., Koyyalamudi, S.R., Pang, G. & et al. (2010). White button mushrooms lowers blood glucose and cholesterol levels in diabetic and hypercholesterolemic rats. *Nutrition Research*, 30, 49–56.
- Jie, J., Zhang, X., Hu, Y.S., Wu, Y., Wang, Q.Z., Li, N.N. & et al. (2009). Evaluation of *in vivo* antioxidant activities of *Ganoderma lucidum* polysaccharides in STZ-diabetic rats. *Food Chemistry*, 115, 32–36.
- Jose, N., Ajith, T.A. & Jananrdhanan, K.K. (2002). Antioxidant, anti-inflammatory and antitumor activities of culinary-medicinal mushroom *Pleurotus pulmonarius* (Fr.) Quel. (Agaricomycetideae). *International Journal of Medicinal Mushrooms*, 4, 329–335.

- Jose, N. & Janardhanan, K.K. (2000). Antioxidant and antitumour activity of *Pleurotus florida*. *Current Science*, 79(7), 941–943.
- Jocken, J.W.E. & Blaak, E.E. (2008). Catecholamine-induced lipolysis in adipose tissue and skeletal muscle in obesity. *Physiology & Behavior*, 94, 219–230.
- Jocken, J.W.E., Blaak, E.E., Kallen, C.J.H., Baak, M.A. & Saris, W.H.M. (2008). Blunted  $\beta$ -adrenoceptor-mediated fat oxidation in overweight subjects: a role for hormone sensitive lipase gene. *Metabolism*, 57, 326–332.
- Jung, S., Choe, J.H., Kim, B., Yun, H., Kruk, Z.A. & Jo, C. (2010). Effect of dietary mixture of gallic acid and linoleic acid on antioxidative potential and quality of breast meat from boilers. *Meat Science*, 86, 520–526.
- Kamat, J.P. & Devasagayam, T.P.A. (1999). Nicotinamide (vitamin B3), as an effective antioxidant against oxidative damage in rat brain mitochondria, *Redox report*, 4(4), 179–184.
- Kanagasabapathy, G., Malek, S.N.A., Kuppusamy, U.R. & Vikineswary, S. (2011). Chemical composition and antioxidant properties of extracts of fresh fruiting bodies of *Pleurotus sajor-caju* (Fr.) Singer. *Journal of Agricultural & Food Chemistry*, 59(6), 2618–2626.
- Kang, S.A., Kim, S.G., Han, Y.Y., Cheong, C., Jang, K.H., Yoon, B.D. & et al. (2008). Inhibitory effect of blood glucose increase by *Aureobasidium* sp. beta-glucan genetic diabetic mice. *Journal of Biotechnology*, 136, 252–260.
- Kang, J., Wang, H.Q. & Chen, R.Y. (2003). Studies on the constituents of the mycelium produced by fermenting culture of *Flammulina velutipes* Singer. *International Journal of Medicinal Mushrooms*, 5, 391–396.
- Kavishree, S., Hemavathy, J., Lokesh, B.R., Shashirekha, M.N. & Rajarathnam, S. (2008). Fat and fatty acids of Indian edible mushrooms. *Food Chemistry*, 106, 597–602.
- Kasaoka, T.N., Takahashi, M., Kim, H. & Ezaki, O. (1999). Up-regulation of liver uncoupling protein-2 mRNA by either fish oil feeding or fibrate administration to mice. *Biochemical and Biophysical Research Communications*, 257, 879–889.

- Kasai, H. (1997). Analysis of a form of oxidative DNA damage, 8-hydroxy-2-deoxyguanosine, as a marker of cellular oxidative stress during carcinogenesis. *Mutation Research*, 3, 147–163.
- Kawai, K., Motosuno, K & Kasai, H. (2006). Detection of 4-oxo-2-hexenal, a novel mutagenic product of lipid peroxidation, in human diet and cooking vapor. *Mutation Research*, 603, 186-192.
- Keefe, J.H. & Bell, D.S.H. (2007). Postprandial hyperglycemia/Hyperlipidemia is a cardiovascular risk factor. *The American Journal of Cardiology*, 100, 899–904.
- Kenneth, G.A. & Clifton, P.G. (2010). New approaches to the pharmacological treatment of obesity. Can they break through the efficacy barrier? *Pharmacology, Biochemistry & Behavior*, 97, 63–83.
- Kern, P.A., Di, G., Gregoria, B., Lu, T., Rassouli, N. & Ranganathan, G. (2001). Adiponectin expression from human adipose tissue: relation to obesity, insulin resistance and TNF- $\alpha$  expression. *Diabetes*, 52, 1779–1785.
- Kho, Y.S., Abdullah, N., Kuppusamy, U.R. & Oh, I.H. (2009). Antioxidant capacity of fresh and processed fruit bodies and mycelium of *Auricularia aricularia-judae* (Fr.) Quél. *Journal of Medicinal Food*, 12(1), 167-174.
- Kiho, T., Yamane, A., Hui, J., Usui, S. & Ukai, S. (1996). Hypoglycemic activity of a polysaccharide (CS-F30) from the cultural mycelium *Cordyceps sinensis* and its effect on glucose metabolism in mouse liver. *Biological & Pharmaceutical Bulletin*, 19, 294–296.
- Kiho, T., Sobue, S. & Ukai, S. (1994). Structural features and hypoglycemic activities of two polysaccharides from a hot-water extract of *Agrocybe cylindracea*. *Carbohydrate Research*, 251, 81–87.
- Kim, S.Y., Kim, J.H., Kim, S.K., Oh, M.J. & Jung, M.Y. (1994). Antioxidant activities of selected oriental herb extracts. *Journal of American Oil Chemical Society*, 71, 633–640.
- Kim, S.O., Yun, S.J., Jung, B., Lee, E.H., Hahm, D.H., Shim, I. & et al. (2004). Hypolipidemic effects of crude extract of adlay seed in obesity rat fed high fat diet: relations of TNF- $\alpha$  and leptin mRNA expressions and serum lipid levels. *Life Sciences*, 75, 1391–1404.

- Kim, M.Y., Seguin, P., Ahn, J.K., Kim, J.J., Chun, C.S., Kim, E.H. & et al. (2008a). Phenolic compound concentration and antioxidant activities of edible and medicinal mushrooms from Korea. *Journal of Agricultural & Food Chemistry*, 56, 7265–7270.
- Kim, H.Y., Jeong, D.M., Jung, H.J., Jung, Y.J., Yokozawa, T. & Choi, J.S. (2008b). Hypolipidemic effects of *Sophora flavescens* and its constituents in poloxamer 407-induced hyperlipidemic and cholesterol fed rats. *Biological Pharmacology Bulletin*, 31, 73–78.
- Kim, Y.J. & Park, T. (2008). Genes are differentially expressed in the epididymal fat of rats rendered obese by a high-fat diet. *Nutrition Research*, 28, 414–422.
- Kim, S.J., Choi, Y., Choi, Y.H. & Park, T. (2011). Obesity activates toll-like receptor-mediated proinflammatory signaling cascades in the adipose tissue of mice. *Journal of Nutritional Biochemistry*, doi:10.1016/j.nutbio.2010.10.012.
- Kircshbaum, B. (2001). Total urine antioxidant capacity. *Clinical Chimica Acta*, 305, 167–173.
- Kochhar, S.P. & Rossell, J.B. (1990). Detection, estimation, and evaluation of antioxidants in food systems. In: Food Antioxidants edited by Hudson, B.J.F.
- Koh, K.K., Han, S.H. & Quon, M.J. (2005). Inflammatory markers and the metabolic syndrome; insight from therapeutic interventions, *Journal of American College of Cardiology*, 46, 1978–1985.
- Koleva, I.I., van Beek, T.A., Linssen, J.P.H., de Groot, A. & Evastatieva, L.N. (2002). Screening of plant extracts for antioxidant activity: A comparative study on three testing methods. *Phytochemistry Analysis*, 13, 8–17.
- Kong, C.S., Kim, J.A. & Kim, S.K. (2009). Anti-obesity effect of sulfated glucosamine by AMPK signal pathway in 3T3-L1 adipocytes. *Food Chemistry & Toxicology*, 47, 2401–2406.
- Koutelidakis, A., Konstantina, A., Serafini, M., Proestos, C., Komaitis, M., Pecorari, M. & et al. (2009). Green tea, white tea and *Pelargonium purpureum* increase the antioxidant capacity of plasma and some organs in mice. *Basic Nutritional Investigations*, 25, 453–458.



- Krzyczkowski, W., Malinowska, E., Suchocki, P., Kleps, J., Olejnik, M. & Herold, F. (2008). Isolation and quantitative determination of ergosterol peroxide in various edible mushrooms species. *Food Chemistry, Food Chemistry*, 113, 351-355.
- Kubo, K. (2004). Anti-diabetic activity present in the fruit body of *Grifola frondosa* (Maitake). *Biology Pharmacological Bulletin*, 17, 1106-1110.
- Kubola, J. & Siriamornpun, S. (2008). Phenolic contents and antioxidant activities of bitter gourd (*Momordica charantia* L.) leaf, stem and fruit fraction extracts *in vitro*. *Food Chemistry*, 110, 881–890.
- Kuhad, A. & Chopra, K. (2009). Attenuation of diabetic nephropathy by tocotrienol; involvement of NFκB signaling pathway, *Life Sciences*, 84, 296–301.
- Kulisic, T., Radonic, A., Katalinic, V. & Milos, M. (2004). Use of different methods for testing antioxidative activity of oregano essential oil. *Food Chemistry*, 85, 633–640.
- Kuo, C.F., Hsieh, C.H. & Lin, W.Y. (2011). Proteomic response of LAP-activated RAW-264.7 macrophages to the anti-inflammatory property of fungal ergosterol. *Food Chemistry*, 126, 207–212.
- Kwiterovich, P.O. (2000). Metabolic pathway of high density lipoprotein and triacylglycerols. *Cardiology*, 86, 120–128.
- Lass, A., Zimmermann, R., Oberer, M. & Zechner, R. (2011). Lipolysis- a highly regulated multi-enzyme complex mediates the catabolism of cellular fat stores. *Progress in Lipid Research*, 50, 14–27.
- Lee, H.S., Park, H.J., Chun, H.K., Cho, S.Y., Cho, S.M. & Lillehoj, H.S. (2006). Dietary phytic acids lowers the blood glucose level in diabetic KK mice. *Nutrition Research*, 26, 474–479.
- Lee, Y.L., Huang, G.W., Liang, Z.C. & Mau, J.L. (2007). Antioxidant properties of three extracts from *Pleurotus citrinopileatus*. *LWT*, 40, 823–833.
- Lee, L., Alloosh, M., Saxena, R., Van Alstine, W., Watkins, B.A., Klaunig, J.E. & et al. (2009). Nutritional model of steatohepatitis and metabolic syndrome in the Ossabaw miniature swine. *Hepatology*, 50, 56–67.

- Lee, H., Kang, R. & Yoon, Y. (2010). SH21B, an anti obesity herbal composition, inhibits fat accumulation in 3T3-L1 adipocytes and high-fat diet induced obese mice through modulation of the adipogenesis pathway. *Journal of Ethnopharmacology*, 127, 709–717.
- Letizia, C.S., Cocchiara, J., Lapczynski, A., Lalko, J. & Api, A.M. (2005). Fragrance material review on cinammic acid. *Food Chemistry & Toxicology*, 43, 925–943.
- Leeuwenburgh, C., Hansen, P.A., Holloszy, J.O. & Heinecke, J.W. (1999). Oxidized amino acids in the urine of aging rats: potential markers for assessing oxidative stress in vivo. *American Journal of Physiology*, 276, R128–R135.
- Li, J., Yu, X., Pan, W. & Unger, R.H. (2002). Gene expression profile of rat adipose tissue the onset of high-fat-diet obesity. *American Journal of Physiology; Endocrinology and Metabolism*, 282, 1334–1341.
- Liao, C.H. & Lin, J.Y. (2011). Lotus plumule polysaccharides protect the spleen and liver from spontaneous inflammation in non-obese diabetic mice by modulating pro-/anti inflammatory cytokine gene expression. *Food Chemistry*, doi:10.1016/j.foodchem.2011.03.103.
- Liu, F.Y., Luo, K.W., Yu, Z.M., Co, N.N., Wu, S.H., Wu, P. & et al. (2009). Suillin from the mushroom *Suillus placidus* as potent apoptosis inducer in human hepatoma HepG2 cells. *Chemico-Biological Interactions*, 181, 168–174.
- Liu, B.H., Wang, Y.C., Wu, S.H., Mersamann, H.J., Cheng, W.T.K. & Ding, S.T. (2008). Insulin regulates the expression of adiponectin and adiponectin receptors in adipocytes. *Domestic Animal Endocrinology*, 34, 352–359.
- Livak, K.J. & Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta C_t}$  method. *Methods*, 25, 402–408.
- Lo, K.M. & Cheung P.C.K. (2004). Antioxidant activity of extracts from the fruiting bodies of *Agrocybe aegerita* var. *alba*. *Food Chemistry*, 89(4), 533–539.
- Lo, H.C., Tu, S.T., Lin, K.C. & Lin, S.C. (2004). The anti-hyperglycemic activity of the fruiting body of *Cordyceps* in diabetic rats induced by nicotinamide and streptozotocin. *Life Sciences*, 74, 2897–2908.

- Lo, S.C., Ho, Y.S. & Buswell, J.A. (2001). Effects of phenolic monomers on the production of laccase by the edible mushrooms *Pleurotus sajor-caju* and, partial characterization of a major laccase component. *Mycologia*, 93(3), 413–421.
- Long, E.K. & Picklo, M.J.S. (2010). Trans-4-hydroxy-2-hexenal, a product of n-3-fatty acid peroxidation: Make some room HNE... *Free Radical Biology*, 49, 1–8.
- Madsen, L., Petersen, R.K., Sorensen, M.B., Jorgensen, C., Hallenborg, P., Pridals, L. & et al. (2003). Adipocyte differentiation of 3T3-L1 preadipocytes is dependent on lipoxygenase activity during the initial stages of the differentiation process. *Biochemistry Journal*, 375, 539–549.
- Mallavadhani, U.V., Sudhakar, A.V.S., Satyanarayana, K.V.S., Mahapatra, A., Li, W. & vanBreemen, R.B. (2004). Chemical and analytical screening of some edible mushrooms. *Food Chemistry*, 95, 58–64.
- Mantovani, M.S., Bellini, M.F., Angeli, J.P.F., Oliveira, R.J., Silva, A.F. & Rebeiro, L.R. (2008). B-Glucans in promoting health: Prevention against mutation and cancer. *Mutation Research*, 658, 15–161.
- Manzi, P. & Pizzoferrato, L., (2000). Beta-glucans in edible mushrooms. *Food Chemistry*, 68, 315–318.
- Manzi, P., Gambelli, L., Marconi, S., Vivanti, V., & Pizzoferrato, L. (1999). Nutrients in edible mushrooms: an interspecies comparative study. *Food Chemistry*, 65(4), 477–482.
- Mathe, D. (1995). Dyslipidemia and diabetes: animal models. *Diabetes Metabolism*, 21, 106–111.
- Mathews, D.R., Hosker, J.P., Rudenski, A.S., Naylor, B.A., Treacher, D.F. & Turner, R.C. (1985). Homeostasis model assessment: insulin resistant and B-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*, 28, 412–419.
- Matilla, P., Konko, K., Eurola, M., Pihlaya, J.M., Astola, J., Vahteristo, L. & et al. (2001). Contents of vitamins, minerals elements and some phenolic compounds in cultivated mushrooms. *Journal of Agricultural & Food Chemistry*, 49, 2343–2348.

- Matsuura, F., Yamashita, S., Nakamura, T., Nishida, M., Nozaki, S., Funahashi, T. & et al. (1998). Effect of visceral fat accumulation on uric acid metabolism in male obese subjects: visceral fat obesity is linked more closely to overproduction of uric acid than subcutaneous fat obesity. *Metabolism*, 47, 929–933.
- Mau, J.L., Chieh, N.C., Shih, J.H. & Chin, C.C. (2004). Antioxidative properties of methanolic extracts from *Grifola frondosa*, *Morchella esculenta* and *Termitomyces albuminosus* mycelia. *Food Chemistry*, 87(1), 111–118.
- Mau, J.L., Lin, H.C. & Song, S.F. (2002). Antioxidant properties of several specialty mushrooms. *Food Research International*, 35, 519–526.
- Maynard, C. (2001). Antioxidants: Food for your body's defense. *Current Health* 2, 27(7), 27–39.
- McChesney, J.D., Venkataraman, S.K. & Henri, J.T. (2007). Plant natural products: Back to the future or into extinction. *Phytochemistry*, 68, 2015–2022.
- Menezes, J.C.J.M.D.S., Kamat, S.P., Cavaleiro, J.A., Gaspar, A., Garrido, J. & Borges, F. (2011). Synthesis and antioxidant activity of long chain alkyl hydroxycinnamates. *European Journal of Medicinal Chemistry*, 46, 773–777.
- Mertens, I.L. & Gaal, V.L.F. (2000). Overweight, obesity, and blood pressure: the effects of modest weight reduction. *Obesity Review*, 8, 270–278.
- Miles, P.G. & Chang, S.T. (1997). *Mushroom Biology-Concise Basics and Current Developments*. World Scientific, USA.
- Mimura, M., Nabeshima, R., Maeda, M. & Shiomi, N. (2009). A highly sensitive enzyme-linked immunosorbent assay for quantification of adipocytokines secreted by mouse adipocytes. *Biochemical Engineering Journal*, 43, 58–63.
- Ming, M., Guanhua, L., Zhanhai, Y., Guang, C. & Xuan, Z. (2009). Effect of the *Lycium barbarum* polysaccharides administration on blood lipid metabolism and oxidative stress of mice fed high-fat diet in vivo. *Food Chemistry*, 113, 872–877.
- Miura, S., Tsunoda, N., Ikeda, S., Kai, Y., Ono, M., Maruyama, K. & et al. (2003). Regulatory sequence elements of mouse GLUT4 gene expression in adipose tissue. *Biochemical & Biophysical Research Communications*, 312, 277–284.

- Miyata, Y., Tanaka, H., Shimada, A., Sato, K., Ito, A., Yamanouchi, T. & et al. (2011). Regulation of adipocytokine secretion and adipocyte hypertrophy by polymethoxyflavonoids, nobiletin and tangeretin. *Life Sciences*, 88, 613–618.
- Mizuno, T. & Zhuang, C. (1995). Houbitake, *Pleurotus sajor-caju*: Antitumor activity and utilization. *Food Reviews International*, 11(1), 185–187.
- Mizuno, T. (1995). Health foods and medicinal usages of mushrooms. *Food Reviews International*, 12, 851–873.
- Mohanty, P., Ghanim, H., Hamouda, W., Aljada, A., Garg, R., Dandona, P. (2002). Both lipid and protein intakes stimulate increased generation of reactive oxygen species by polymorphonuclear leukocytes and mononuclear cells. *American Journal of Clinical Nutraceuticals*, 75, 767–772.
- Mohanty, P., Aljada, A., Ghanim, H., Hofmeyer, D., Tripathy, D., Syed, T. & et al. (2004). Evidence for a potent anti-inflammatory effect of rosiglitazone. *Journal of Clinical Endocrinology Metabolism*, 89, 2728–2735.
- Moon, H.S., Lee, H.G., Seo, G.H., Chung, C.H. Kim, T.G., Choi, Y.J. & et al. (2009). Antiobesity effect of PEGylated conjugated linoleic acid on high fat diet induced obese mice C57BL/6J (*ob/ob*) mice: attenuation of insulin resistance and enhancement of antioxidant defenses. *Journal of Nutritional Biochemistry*, 20, 187–194.
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunology Methods*, 65(1), 55–63.
- Moskovitz, J., Yim, B.M. & Chock, P.B. (2002). Free radicals and disease. *Archives of Biochemistry & Biophysics*, 397, 354–359.
- Mukhopadhyay, R., Chatterjee, S., Chatterjee, B.P. & Guha, A.K. (2005). Enhancement of biomass production of edible mushroom *Pleurotus sajor-caju* grown in whey by plant growth hormones. *Process Biochemistry*, 40, 1241–1244.
- Muller, S., Martin, S., Koenig, W., Hanifi-Moghaddam, P., Rathmann, W., Haastert, B. & et al. (2002). Impaired glucose tolerance is associated with increased serum concentrations of interleukin 6 and co-regulated acute-phase proteins but not TNF- $\alpha$  or its receptors. *Diabetologia*, 45, 805–812.

- Nagata, M., Ijichi, S., Hashiguchi, T., Morioka, H. & Arimura, K. (2009). Metabolic syndrome-related markers in obese university students in Kagoshima, Japan: Implications of uric acid and plasminogen activator inhibitor type-1. *Diabetes & Metabolic syndrome: Clinical research & Reviews*, 3, 96–102.
- Nagata, M., Takamura, T., Ando, H., Nakamura, S., Kurita, S., Misu, H. & et al. (2008). Increased oxidative stress precedes the onset of high-fat diet induced insulin resistance and obesity. *Metabolism Clinical and Experimental*, 57, 1071–1077.
- Ng, T.B., Lam, S.K. & Chan, S.Y. (2002). A ubiquitin-like peptide from the mushroom *Pleurotus sajor-caju* exhibits relatively potent translation-inhibitory and ribonuclease activities. *Peptides*, 23, 1361–1365.
- Ngai, P.H.K. & Ng, T.B. (2004). A ribonuclease with microbial, antimitogenic and antiproliferative activities from the edible mushroom *Pleurotus sajor-caju*. *Peptides*, 25, 11–17.
- Ngai, P.H.K. & Ng, T.B. (2003). Lentin, a novel and potent antifungal protein from shitake mushroom with inhibitory effects on activity of human immunodeficiency virus-1 reverse transcriptase and proliferation of leukemia cells. *Life Sciences*, 73, 3363–3374.
- Newman, D.J., Cragg, G.M. & Snader, K.M. (2003). Natural product as sources of new drugs over the period 1981-2002. *Journal of Natural Products*, 66, 1022–1037.
- Nieto, I.J. & Chegwin, A.C. (2008). Triterpenoids and fatty acids identified in the edible mushroom *Pleurotus sajor-caju*. *Journal of Chilean Chemical Society*, 53, 1515–1517.
- Niki, E. (2010). Assessment of antioxidant capacity *in vitro* and *in vivo*. *Free Radical Biology & Medicine*, 49, 505–515.
- Ohkoshi, E., Miyazaki, H., Shindo, K., Watanabe, H., Yoshida, A. & Yajima, H. (2007). Constituents from the leaves of *Nelumbo nucifera* stimulate lipolysis in the white adipose tissue of mice. *Planta Medica*, 73(12), 1255–1259.
- Ohshima, H. & Bartsch, H. (1994). Chronic infections and inflammatory processes as cancer risk factors: possible role of nitric oxide in carcinogenesis. *Mutation Research*, 305, 253–264.

- Ono, Y., Hattori, E., Fukaya, Y., Imai S. & Ohizumi, S. (2006). Anti-obesity effect of *Nelumbo nucifera* leaves extract in mice and rats. *Journal of Ethnopharmacology*, 106, 238–244.
- Orhan, I & Ustun, O. (2011). Determination of total phenol content, antioxidant activity and acetylcholinesterase inhibition in selected mushrooms from Turkey. *Journal of Food Composition & Analysis*, doi:10.1016/j.jfca.2010.11.005.
- Oyetayo, V.O. (2008). Mushrooms Indigenous to Nigeria as potential source of myconutraceuticals- A review. *Current Trends in Biotechnology & Pharmacology*, 2(4), 471–477.
- Pai, S.H., Jong, S.C. & Low, D.W. (1990). Usages of mushroom. *Bioindustry*, 1, 126–131.
- Palacios, I., Lozano, M., Moro, C., D'Arrigo, M., Rostagno, M.A. & Martinez, J.A. (2011). Antioxidant properties of phenolic compounds occurring in edible mushrooms. *Food Chemistry*, 128, 674–678.
- Pannacciulli, N., Cantatore, F.P., Minenna, A., Bellacicco, M., Giorgino, R. & De Pergola, G. (2001). C-reactive protein is independently associated with total body fat, central fat, and insulin resistance in adult women. *International Journal of Obesity Related Metabolism Disorder*, 25, 1416–1420.
- Pasimeni, G., Ribaudo, M.C., Capoccia, D., Rossi, F., Bertone, C., Leonetti, F. & et al. (2006). Non invasive evaluation of endothelial dysfunction in uncomplicated obesity: relationship with insulin resistance. *Microvascular Research*, 71, 115–120.
- Park, S.H., Ko, S.K. & Chung, S.H. (2005). *Euonymus alatus* prevents hyperglycemia and hyperlipidemia induced by high-fat diet in ICR mice. *Journal of Ethnopharmacology*, 102, 326–335.
- Pramanik, M., Mondal, S., Chakraborty, I., Rout, D. & Islam, S.S. (2005). Structural investigation of polysaccharides (Fr II) isolated from aqueous extract of an edible mushroom, *Pleurotus sajor-caju*. *Carbohydrate Research*, 340, 629–636.
- Pramanik, M., Mondal, S., Chakraborty, I., Rout, D. & Islam, S.S. (2007). Structural analysis of a water-soluble glucan (Fr I) of an edible mushroom, *Pleurotus sajor-caju*. *Carbohydrate Research*, 342, 2670–2675.

- Pellegrini, N., Serafini, M., Colombi, B., Del Rio, D., Salvatore, S., Bianchi, M. & et al. (2003). Total antioxidant capacity of plant foods, beverages and oils consumed in Italy assessed by three different *in vitro* assays. *Nutrient Requirements*, 133, 2812–2819.
- Peng, Y., Rideout, D., Rakita, S., Sajan, M., Farese, R., You, M. & et al. (2009). Downregulation of adiponectin/adipor2 is associated with steatohepatitis in obese mice. *Journal of Gastrointestinal Surgery*, 13, 2043–2049.
- Perera, C.O., Jayasinghe, V.J., Ng, F.L. & Mujumdar, A.S. (2003). The effects of moisture content on the conversion of ergosterol to vitamin D in shiitake mushrooms. *Drying Technology*, 21(6), 1091–1099.
- Phillips, M.S., Liu, Q., Hammod, H.A., Dugan, V., Hey, P.J., Caskey, C.J. & et al. (1996). Leptin receptor missense mutation in the fatty Zucker rat. *Natural Genetics*, 13, 18–19.
- Phillipson, J.D. (2001). Phytochemistry and medicinal plants. *Phytochemistry*, 56, 237–243.
- Pilch, P.F. & Bergenheim, N. (2006). Pharmacological targeting of adipocytes/fat metabolism for treatment of obesity and diabetes. *Molecular Pharmacology*, 70, 779–785.
- Poulos, S.P., Dodson, M.V. & Hausman, G.J. (2010). Cell lines models for differentiation: preadipocytes and adipocytes. *Experimental Biology & Medicine*, 235, 1185–1193.
- Puente, B.A., Fève, B., Fellahi, S. & Bastard, J.P. (2008). Adipokines: the missing link between insulin resistance and obesity. *Diabetes & Metabolism*, 34, 2–11.
- Quimio, T.H. (2001). Edible mushroom: Potential and Low Cost Technologies for Tropical Cultivation in Trichomycetes and Other Fungal Groups. Science Publishers Inc. USA.
- Rains, J.L. & Jain, S.K. (2011). Oxidative stress, insulin signaling and diabetes. *Free Radical Biology & Medicine*, 50, 567–575.



- Ramarathnam, N., Osawa, T., Ochi, H. & Kawakishi, S. (1995). The contribution of plant food antioxidants to human health. *Trends in Food Science & Technology*, 6, 75–82.
- Reaven, G. (1988). Role of insulin resistance in human disease. *Diabetes*, 37, 1595–1607.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M. & Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology & Medicine*, 26, 1231–1237.
- Reich, H.J., Jautelat, M., Messe, M.T., Weigert, F.J. & Roberts, J.D. (1969). Nuclear magnetic resonance spectroscopy. Carbon 13 spectra of steroids. *Journal of American Chemical Society*, 91, 7445–7454.
- Rio, D.D., Stewart, A.J. & Pellegrini, N. (2005). A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutrition Metabolism & Cardiovascular Diseases*, 4, 216–328.
- Robidoux, J., Martin, T.L. & Collins, S. (2004).  $\beta$ -adrenergic receptors and regulation of energy expenditure: a family affair. *Annual Review of Chemistry*, 253, 7570–7578.
- Roberts, C.K. & Sindhu, K.K. (2009). Oxidative stress and metabolic syndrome. *Life Sciences*, 84, 705–712.
- Rosen, G.M., Pou, S., Ramos, C.L., Cohen, M.S. & Britigan, B.E. (1995). Free radicals and phagocytic cells. *FASEB Journal*, 9, 200–209.
- Roy, S.K., Maiti, D., Mondal, S., Das, D. & Islam, S.S. (2008). Structural analysis of a polysaccharide isolated from the aqueous extract of an edible mushroom, *Pleurotus sajor-caju*, cultivar Black Japan. *Carbohydrate Research*, 343, 1108–1113.
- Rudich, A., Tirosh, A., Potashnik, R., Khamaisi, N. & Bashan, N. (1999). Lipoic acid protects against oxidative stress induced impairment in insulin stimulation of protein kinase B and glucose transport in 3T3-L1 adipocytes. *Diabetologia*, 42, 949–957.
- Ruthig, D.J. & Meckling-Gill, K.A. (1999). Both (n-3) and (n-6) fatty acids stimulate wound healing in the rat intestinal epithelial cell line, IEC-6. *Journal of Nutrition*, 129 (10), 1791–1798.

- Sakanaka, S., Tachibana, Y., Ishihara, N. & Juneja, L.R. (2004). Antioxidants activity of egg-yolk protein hydrolysates in a linoleic acid oxidation system. *Food Chemistry*, 86(1), 99–103.
- Said, S., Neves, F.M., & Griffiths, A.J.F. (2004). Cinnamic acids inhibits the growth of the fungus *Neurospora crassa*, but is eliminated as acetophenone. *International Biodeterioration & Biodegradation*, 54, 1–6.
- Saito, T., Abe, D. & Sekiya, K. (2007). Nobiletin enhances differentiation and lipolysis of 3T3-L1 adipocytes. *Biochemical & Biophysical Research Communication*, 357, 371–376.
- Saltarelli, R., Ceccaroli, P., Iotti, M., Zamnonelli, A., Buffalini, M., Casadei, L. & et al. (2009). Biochemical characterization and antioxidant activity of mycelium of *Ganoderma lucidum* from Central Italy, *Food Chemistry*, 116, 143–151.
- Saito, F., Iwamoto, S. & Yamauchi, R. (2011). 4-oxo-2-nonanal as a pro-oxidant during autoxidation of methyl linoleate in bulk phase. *Food Chemistry*, 124, 1496–1499.
- Sakurai, T., Kitadate, K., Nishioka, H., Fujii, H., Kizaki, T., Kondoh, Y. & et al. (2010). Oligomerized grape seed polyphenols attenuate inflammatory changes due to antioxidative properties in coculture of adipocytes and macrophages. *Journal of Nutrition & Biochemistry*, 21, 47–54.
- Sanchez, D., Quinones, M., Moulay, L., Muguerza, B., Miguel, M. & Aleixandre, A. (2011). Soluble fiber-enriched diets improve inflammation and oxidative stress biomarkers in Zucker fatty rats. *Pharmacological Research*, doi:10.1016/j.phrs.2011.02.005.
- Sarikurkcü, C., Tepe, B. & Yamac, M. (2008). Evaluation of the antioxidant activity of four edible mushrooms from central Anatolia, Eskisehir-Turkey: *Lactarius deterrimus*, *Suillus collitinus*, *Boletus edulis*, *Xerocomus chrysenteron*. *Bioresource Technology*, 99, 6651–6655.
- Sarikurkcü, C., Tepe, B., Semiz, D.K. & Solak, M.H. (2010). Evaluation of metal concentration and antioxidant activity of three edible mushrooms from Mugla, Turkey. *Food Chemical Toxicology*, 48, 1230–1233.
- Sarmadi, B.H. & Ismail, A. (2010). Antioxidative peptides from food proteins: A review. *Peptides*, 31, 1949–1956.

- Sauve, A.A. & Xu, P. (2010). Vitamin B3, the nicotinamide adenine dinucleotides and aging. *Mechanism & Ageing Development*, 131, 287–298.
- Scheja, L., Heese, B. & Seedorf, K. (2011). Beneficial effects of IKK –deficiency on body weight and insulin sensitivity are lost in high fat diet-induced obesity in mice. *Biochemical & Biophysical Research Communications*, 407, 288–294.
- Schreyer, S.A., Wilson, D.L., & LeBoeuf, C.F. (1998). C57BL/6 mice fed high fat diets as model for diabetes accelerated atherosclerosis. *Atherosclerosis*, 136, 17–24.
- Schleiser, K., Harwat, M., Bohm, V. & Bitsch, R. (2002). Assessment of antioxidant activity by using different in vitro methods. *Free Radical Research*, 36, 177–187.
- Schneider, I., Kressel, G., Meyer, A., Krings, U., Berger, R.G. & Hahn, A. (2011). Lipid lowering effects of oyster mushroom (*Pleurotus ostreatus*) in humans. *Journal of Functional Foods*, 3, 17–24.
- Seto, S.W., Lam, T.Y., Tam, H.L., Au, A.L.S., Chan, S.W., Wu, J.H. & et al. (2009). Novel hypoglycemic effects of *Ganoderma lucidum* water-extract in obese/diabetic (+db/+db) mice. *Phytomedicine*, 16, 426–436.
- Shamtsyan, M., Konusova, V., Maksimova, Y., Goloshchev, A., Panchenko, A., Simbirtsev, A. & et al. (2004). Immunomodulating and anti-tumor action of extracts of several mushrooms. *Journal of Biotechnology*, 113, 77-83.
- Shi, Y., Chen, Q.X., Wang, Q., Song, K.K. & Qui, L. (2005). Inhibitory effects of cinnamic acid and its derivatives on the diphenolase activity of mushroom (*Agaricus bisporus*) tyrosinase. *Food Chemistry*, 92, 707–712.
- Shu, Y.Z. (1998). Recent natural products based drug development: A pharmaceutical industry perspective. *Journal of Natural Products*, 61, 1053–1071.
- Silva, L., Garcia, B. & Martins, F.P. (2010). Oxidative stability of olive oil and polyphenolic compounds after boiling vegetable process. *LWT-Food Science Technology*, 43, 1336–1344.
- Sim, K.S., Nurestri, S.A.M. & Norhanom, A.W. (2010). Phenolic content and antioxidant activity of crude and fractionated extracts of *Pereskia bleo* (Kunth) DC. *African Journal of Pharmacy & Pharmacology*, 4(5), 193–201.

- Slovacek, L., Pavlik, V. & Slovackova, B. (2008). The effect of sibutramine therapy on occurrence of depression symptoms among obese patients. *Nutrition, Metabolism & Cardiovascular Disease*, 18, 43–44.
- Shoelson, S.E., Laura, H. & Naaz, A. (2007). Obesity, inflammation and insulin resistance. *Gastroenterology*, 132, 2169–2180.
- Smiderle, F.R., Olsen, L.M., Carbonero, E.R., Baggio, C.H., Freitas, C.S., Marcon, R. & et al. (2008). Anti-inflammatory and analgesic properties in a rodent model of a (1→3)(1→6)-linked  $\beta$ -glucan isolated from *Pleurotus pulmonarius*. *European Journal of Pharmacology*, 597, 86–91.
- Smith, U., Andersson, C.X., Gustafson, B., Hammarstedt, A., Isakson, P. & Wallerstedt, E. (2007). Adipokines, systemic inflammation and inflamed adipose tissue in obesity and insulin resistance. *International Congress Series*, 1303, 31–34.
- Soares, A.F., Guichardant, M., Cozzone, D., Hubac, N.B., Tiali, N.B. & Geloën, L.A. (2005). Effects of oxidative stress on adiponectin secretion and lactate production in 3T3-L1 adipocytes. *Free radical Biology & Medicine*, 38, 882–889.
- Song, Y.S., Kim, S.H., Sa, J.H., Jin, C., Lim, C.J. & Park, E.H. (2003). Anti-angiogenic, antioxidant and xanthine oxidase inhibition activities of mushroom *Phellinus linteus*. *Journal of Ethnopharmacology*, 88, 113–116.
- Soobratte, M.A., Neergheen, V.S., Baborun, T. & Luximon-Ramma, O.I.A. (2005). Phenolics as potential antioxidant therapeutic agents: Mechanism and actions. *Fundamental & Molecular Mechanisms of Mutagenesis*, 579, 200–213.
- Soong, Y.Y. & Barlow, P.J. (2004). Antioxidant activity and phenolic content of selected fruits seeds. *Food Chemistry*, 88, 411–417.
- Spiegelman, B.M. & Flier, J.S. (2001). Obesity and regulation of energy balance. *Cell*, 104, 531–534.
- Sri Nurestri A.M., Norhanum A. W., Hashim Y., Sim, K.S., Hong, S.L., Lee, G.S. & et al. (2008). Cytotoxic activity of *Pereskia bleo* (Cactaceae) against selected human cell lines. *International Journal of Cancer Research*, 4(1), 20–27.

- Stamets, P. (2005). Mycelium running-How mushrooms can help save the world. Ten Speed Press, California, USA, 202–205
- Stern, K.R. (2006). Introductory Plant Biology. (Ed: 10<sup>th</sup>). Mc Graw Hill Companies, Inc. USA, 245–250
- Stewart, L.K., Soileu, J.L., Ribnicky, D., Wang, Z.Q., Raskin, I., Poulev, A. & et al. (2008). Quercetin transiently increases energy expenditure but persistently decreases circulating markers of inflammation in C57BL/6J mice fed high –fat diet. *Metabolism Clinical & Experimental*, 57, 39–46.
- Su, C.L., Sztalryd, C., Contreras, J.A., Holm, C., Kimmel, A.R. & Londos, C. (2003). Mutational analysis of the hormone sensitive lipase translocation reaction in adipocytes. *Journal of Biological Chemistry*, 278, 43615–43619.
- Subauste, A.R. & Burant, C.F. (2007). Role of FOXO1 in FFA –induced oxidative stress in adipocytes. *American Journal of Physiology Endocrinology Metabolism*, 293, 159–164.
- Suja, K.P., Jayalekshmy, A. & Arumughan, C. (2005). Antioxidant activity of sesame cake extract. *Food Chemistry*, 91, 213–219
- Sun, J.E., Ao, Z.H., Lu, Z.M., Xu, H.Y., Zhang, X.M., Dou, W.F. & et al. (2008). Antihyperglycemic and antilipidprotective effects of dry matter of culture broth of *Inonotus obliquus* in submerged culture on normal and alloxan-diabetes mice. *Journal of Ethnopharmacology*, 118, 7–13.
- Surwit, R.S., Feinglos, M.N., Rodin, J., Sutherland, A., Petro, A.E., Opara, A.C. & et al. (1995). Differential effects of fat and sucrose on the development of obesity and diabetes in C57BL6J mice. *Metabolism*, 44, 645–651.
- Szkuldeski, T., Nogowski, L., Oszmalek, E.W., Kaczmarek, P. & Szkuldeski, K. (2005). Genistein restricts leptin secretion from rat adipocytes. *Journal of Steroid Biochemistry & Molecular Biology*, 96, 301–306.
- Szkuldelski, K., Szkuldelski, T. & Nogowski, L. (2002). Daidzein, coumestrol and zearalenone affect lipolysis in rat adipocytes. *Phytomedicine*, 9, 338–345.

- Takaku, T., Kimura, Y. & Okuda, H. (2001). Isolation of an antitumor compound from *Agaricus blazei* Murill and its mechanism of action. *Biochemical & Molecular of Nutrients*, 1409–1413.
- Talior, I., Tennenbaum, T., Kuroki, T. & Finkelman, H.E. (2005). PKC-D- dependent activation of oxidative stress in adipocytes of obese and insulin resistant mice: role for NADPH oxidase. *American Journal Physiology Endocrinology & Metabolism*, 288, 405–411.
- Tan, M.J., Turner, N., Ye, J.M., Behrens, H.C., Ke, C.Q., Tang, C.P. & Ye, Y. (2008). Antidiabetic activities of triterpenoids isolated from bitter melon associated with activation of the AMPK pathway. *Journal of Chemical Biology*, 15, 263–273.
- Telles, C.B.S., Sabry, D.A., Lima, J.A., Costa, M.S.S.P., Silveira, R.F.M., Trindade, E.S. & et al. (2011). Sulfation of the extracellular polysaccharide produced by the edible mushroom *Pleurotus sajor-caju* alters its antioxidant, anticoagulant and antiproliferative properties *in vitro*. *Carbohydrate Polymers*, 85, 514–521
- Terra, X., Pallares, V., Ardevol, A., Blade, C., Larrea, J.F., Pujadas, G. & et al. (2011). Modulatory effect of grape seed procyanidins on local systemic inflammation in diet induced obesity rats. *Journal of Nutritional Biochemistry*, 22, 380–387.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L. & Byrne, D.H. (2006). Comparison of ABTS, DPPH, FRAP and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition & Analysis*, 19, 669–675.
- Thetsrimuang, C., Khammuang, S. & Samthima, S. (2011). Antioxidant activity of crude polysaccharides from edible fresh and dry mushroom fruiting bodies of *Lentinus* sp. strain RJ-2. *International Journal of Pharmacology*, 7 (1), 58–65.
- Tilg, H. & Moschen, A.R. (2006). Adipocytokines: mediators linking adipose tissue, inflammation and immunity. *Natural Review Immunology*, 6, 772–783.
- Tsai, S.Y., Huang, S.J., Lo, S.H., Wu, T.P., Lian, P.Y. & Mau, J.L. (2009). Flavour components and antioxidant properties of several cultivated mushrooms. *Food Chemistry*, 113, 578–584.

- Tsigos, C., Papanicolaou, D.A., Kyrou, I., Defensor, R., Mitsiadis, C.S. & Chrousos, G.P. (1997). Dose-dependent effects of recombinant human interleukin-6 on glucose regulation. *Journal of Endocrinology Metabolism*, 82, 4167–4170.
- Tsuchiya, K., Sakai, H., Suzuki, N., Iwashima, F., Yoshimoto, T., Shichiri, M. & et al. (2007). Chronic blockade of nitric oxide synthesis reduces adiposity and improves insulin resistance in high fat-induced obese mice. *Endocrinology*, 148, 4548–4556.
- Townsend, K.L., Lorenzi, M.M. & Widmaier, E.P. (2008). High-fat diet-induced changes in body mass and hypothalamic gene expression in wild-type and leptin-deficient mice. *Endocrine*, 33, 176–188.
- Toyosaki, T., Sakane, Y. & Kasai, M. (2008). Oxidative stability, trans -2,4-decadienals and tocopherol contents during storage of dough fried in soybean oil with added medium-chain triacylglycerols. *Food Research International*, 41, 318–324.
- Turner, L. (1997). Antioxidants increase our odds for achieving optimal health. *Better Nutrition*, 59(7), 22–30.
- Ukawa, Y., Ito, H. & Hisamatsu, M. (2000). Antitumor effects of (1→3)-β-D-glucan and (1→6)-β-D-glucan purified from newly cultivated mushroom, Hatakeshimeji (*Lyophyllum decastes* Sing). *Journal of Bioscience & Bioengineering*, 90(1), 98–104.
- Umar, A.R., Ahmed, Q.U., Muhammad, B.Y., Dogarai, B.B.S. & Mat Soad, S.Z. (2010). Anti-hyperglycemic activity of the leaves of *Tetracera scandens* Linn. Merr. in allaxon induced diabetic rats. *Journal of Ethnopharmacology*, 131, 140–145.
- Vaz, J.A., Barros, L., Martins, A., Santos-Buelga, C., Vasconcelos, M.H. & Ferreira, I.C.F.R. (2010). Chemical composition of wild edible mushrooms and antioxidant properties of their water soluble polysaccharidic and ethanolic fractions. *Food Chemistry*, 126, 610–616.
- Vaz, J.A., Heleno, S.A., Martins, A., Almeida, G.M., Vasconcelos, M.H. & Ferreira, I.C.F.R. (2011). Wild mushrooms *Clitocybe alexandri* and *Lepista inversa*: *In vitro* antioxidant activity and growth inhibition of human tumour cell lines. *Food Chemistry Toxicology*, 48, 2881–2884.
- Vroegrijk, I.C.O.M., Diepan, J.A.V., Berg, S.V.D., Weatbroek, I., Keizer, H., Gambelli, L. & et al. (2011). Pomegranate seed oil, a rich source of puninic acid, prevents diet-

induced obesity and insulin resistance in mice. *Food & Chemical Toxicology*, doi:10.1016/j.fct.2011.03.037.

Wannamethee, S.G. & Shaper, A.G. (1999). Weight change and duration of overweight and obesity in the incidence of type 2 diabetes. *Diabetes Care*, 22, 1266–1272.

Wang, Y., Barbacioru, C., Hyland, F., Xiao, W., Hunkapiller, K.L., Blake, J. & et al. (2006). Large scale real time PCR validations of gene expression measurements from two commercial long oligonucleotide microarrays. *BMC Genomics*, 7, 59.

Wang, C., Mao, X., Wang, L., Liu, M., Wetzel, M.D. Guan, K.L. & et al. (2007). Adiponectin sensitizes insulin signaling by reducing p70 S6 kinase-mediated serine phosphorylation of IRS-1. *Journal of Biology Chemistry*, 282, 1991–1996.

Wang, S.L., Li, Y., Wen, Y., Chen, Y.F., Na, L.X., Li, S.T. & et al. (2009). Curcumin, a potential inhibitor of up-regulation of TNF- $\alpha$  and IL-6 induced by palmitate in 3T3-L1 adipocytes through NF-kappaB and JNK pathway. *Biomedical Environmental Sciences*, 22, 32–39.

Wasser, S.P. (2002). Medicinal mushrooms as a source of antitumor and immunomodulating polysaccharides. *Applied Microbiology & Biotechnology*, 60, 258–274.

Wasser, S.P., Nevo, E., Sokolov, D., Reshtnikov, S. & Tismenestky, M.T. (2000). Dietary supplements from medicinal mushrooms: Diversity of types and variety of regulations. *International Journal of Medicinal Mushrooms*, 2, 1–19.

Wasser, S.P. & Weis, A.L. (1999a). Medicinal properties of substances occurring in higher Basidiomycetes mushrooms: Current perspectives (Review). *International Journal of Medicinal Mushrooms*, 1, 31–62.

Wasser, S.P. & Weiss, A.L. (1999b). Therapeutics effects of substances occurring in higher Basidiomycetes mushrooms: a modern prospective. *Critical Review on Immunology*, 19(1), 65–96.

Wasser, S.P. & Weis, A.L. (1999c). General description of the most important medicinal higher Basidiomycetes mushrooms. *International Journal of Medicinal Mushrooms*, 1, 351–370.



- Wetzel, S., Lachance, H. & Waldmann, H. (2010). Natural products as lead sources for drug development. *Comprehensive Natural Products II*, 3, 5–46.
- Witko-Sarsat, V., Friedlander, M., Khoa, T.N., Blandin, C.C., Nguyen, A., Canteloup, S. & et al. (1998). Advanced oxidation protein products as novel mediators of inflammation and monocyte activation in chronic renal failure. *Journal of Immunology*, 161(5), 2524–2532.
- Wong, K.H., Sabaratnam, V., Abdullah, N., Kuppusamy, U.R. & Naidu, M. (2009). Effects of cultivation techniques and processing on antimicrobial and antioxidant activities on *Hericium erinaceus* (Bull.: Fr.) Pers. extracts. *Food Technology & Biotechnology*, 47(1), 47–55.
- Wood, L.G., Gibson, P.G. & Garg, M.L. (2006). A review of the methodology for assessing in vivo antioxidant capacity. *Journal of Science & Food Agriculture*, 86, 2057–2066.
- Wootton, P.C. & Ryan, L. (2011). Improving public health?: The role of antioxidant rich fruit and vegetables beverages. *Food Research International*, 44, 3135–3148.
- Wu, L.L., Chiou, C.C., Chang, P.Y. & Wu, J.T. (2004). Urinary 8-OHdG: a marker of oxidative stress to DNA and a risk factor for cancer, atherosclerosis and diabetics. *Clinica Chimica Acta*, 1-2, 1–9.
- Xie, G., Schepetkin, A., Siemsen, D.W., Kirpotina, L.N., Wiley, J.A. & Quinn, M.T. (2008). Fractionation and characterization of biologically-active polysaccharides from *Artemisia tripartite*. *Phytochemistry*, 69, 1359–1371.
- Xie, J.T., Wang, C.Z., Li, X.L., Ni, M., Fishbein, A. & Yuan, C.S. (2009). Anti-diabetic effect of American ginseng may not be linked to antioxidant activity: comparison between American ginseng and *Scutellaria baicalensis* using *ob/ob* mice model. *Fitoterapia*, 80, 306–311.
- Yang, R.L., Li, W., Shi, Y.H. & Le, G.W. (2008). Lipoic acid prevents high-fat diet-induced dyslipidemia and oxidative stress: A microarray analysis. *Basic Nutritional Investigations*, 24, 582–588.
- Yang, R.Z., Lee, M.J., Hu, H., Pollin, T.I., Ryan, A.S, Nicklas, B.J. & et al. (2006). Acute-phase serum amyloid A: an inflammatory adipokine and potential link between obesity and its metabolic complications. *PLoS Medicine*, 3, e287.

- Yang, B.K., Kim, D.H., Jeong, S.C., Das, S., Choi, Y.S., Shin, J.S., Lee, S.C. & Song, C.H. (2002). Hypoglycemic effect of a *Lentinus edodes* exo-polymer produced from a submerged mycelial culture. *Bioscience, Biotechnology & Biochemistry*, 66, 937–942.
- Yap, A.T. & Ng, M.L.M. (2001). An improved method for the isolation of lentinan from the edible and medicinal shiitake mushroom (*Lentinus edodes* (Berk.) Sing.). *International Journal of Medicinal Mushrooms*, 3, 6–19.
- Ye, M., Liu, J.K., Lu, Z.X., Zhao, Y., Liu, S.F., Li, L.L. & et al. (2005). Grifolin, a potential antitumor natural product from the mushroom *Albatrellus confluens*, inhibits tumor cell growth by inducing apoptosis in vitro. *FEBS Letters*, 579, 3437–3443.
- Yen, P.L., Chen, B.H., Yang, F.L & Lu, Y.F. (2010). Effects of deep frying oils on blood pressure and oxidative stress in spontaneously hypertensive and normotensive rats. *Nutrition*, 26, 331–336.
- Yin, W., Mu, J. & Birnbaum, M.J. (2003). Role of AMP –activated protein kinase in cyclic AMP- dependent lipolysis in 3T3-L1 adipocytes. *Journal of Biological Chemistry*, 278, 43074–43080.
- Yokota, T., Nagashima, M., Ghazizadeh, M. & Kawanami, O. (2009). Increased effect of fucoidan on lipoprotein lipase secretion in adipocytes. *Life Sciences*, 84, 523–529.
- Yoshino, G., Tanaka, M., Nakano, S., Matsumoto, T., Kojima, M., Murakami, E. & et al. (2009). Effect of Rosvastatin on concentration of plasma lipids, urine and plasma oxidative stress markers, and plasma high-sensitivity c-reactive proteins in hypercholesterolemic patients. *Current Therapeutic Research*, 6, 439–448.
- Yun, J.W. (2010). Possible anti-obesity therapeutics from nature- A review. *Phytochemistry*, 71, 1625–1641.
- Yu, Z.H., Hua, L.Y., Qian, Y. & Yan, L. (2009). Effect of *Lentinus edodes* polysaccharides on oxidative stress, immunity activity and oral ulceration of rats stimulated by phenol. *Carbohydrate Polymers*, 75, 115–118.
- Yuan, Z., He, P., Cui, J. & Takeuchi, H. (1998). Hypoglycemic effect of water-soluble polysaccharide from *Auricularia auricular-judae* Quel. on genetically diabetic KK-Ay mice. *Bioscience, Biotechnology & Biochemistry*, 62, 1898–1903.

- Yuan, J.P., Wang, J.H., Liu, X., Kuang, H.C. & Zhao, S.Y. (2007). Simultaneous determination of free ergosterol and ergosteryl esters in *Cordyceps sinensis* by HPLC. *Food Chemistry*, 105, 1755–1759.
- Yumuk, V.D., Hatemi, H., Tarakchi, T., Uyar, N., Turan, N., Bagriacik, N. & et al. (2005). High prevalence of obesity and diabetes mellitus in Konya, a central Anatolian city in Turkey. *Diabetes Research & Clinical Practice*, 70, 151–158.
- Zhang, W., Liu, C.Q., Wang, P.W., Sun, S.Y., Su, W.J., Zhang, H.J. & et al. (2010). Puerarin improves insulin resistance and modulates adipokine expression in rats fed a high-fat diet. *European Journal of Pharmacology*, 649, 398–402.
- Zhang, M., Cui, S.W., Cheung, P.C.K. & Wang, Q. (2007). Antitumor polysaccharides from mushrooms: a review on their isolation process, structural characteristics and antitumor activity. *Food science & Technology*, 18, 4–19.
- Zhang, M., Cheung, P.C.K., Zhang, L., Chi, M.C. & Cooi, V.E.C. (2004). Carboxymethylated  $\beta$ -glucans from mushroom sclerotium of *Pleurotus tuber-regium* as novel water-soluble anti-tumor agent. *Carbohydrate Polymers*, 57(3), 319–325.
- Zhou, J., Ge, M., Wang, Y., Wu, H., Shen, J. & Ma, X. (2007). Effects of pioglitazone on proliferation and differentiation of human preadipocytes. *Journal of Nanjing Medical University*, 21(1), 99–103.
- Zhuang, C., Mizuno, T., Shimada, A., Ito, H., Suzuki, C., Mayuzumi, Y. & et al. (1993). Antitumor protein-containing polysaccharides from a Chinese mushroom Fengweigu or Houbitake, *Pleurotus sajor-caju* (Fr.) Sings. *Bioscience, Biotechnology & Biochemistry*, 57(6), 901–906.
- Zeigerer, A., Rodeheffer, M.S., McGraw, T.E., Friedman, J.M. (2008). Insulin regulates leptin secretion from 3T3-L1 adipocytes by a P13-kinase independent mechanism. *Experimental Cell Research*, 314, 2249–2256.
- Zou, C. & Shao, J. (2008). Role of adipocytokines in obesity-associated insulin resistance. *Journal of Nutrition Biochemistry*, 19, 277–286.