

# LIST OF FIGURES

---

	<b>Page</b>
<b>Figure 3.1.1:</b> Fractionation and purification of constituents from the leaves of <i>Phyllagathis rotundifolia</i> . M–MCI gel CHP 20P; O–Chromatorex ODS; S–Sephadex LH-20; T–Toyopearl HW-40F; C–Semi-preparative reversed-phase Symmetry C-18 column; F–Fraction.	<b>66</b>
<b>Figure 3.1.2:</b> Fractionation and purification of constituents from the leaves of <i>Phyllagathis praetermissa</i> . M–MCI gel CHP 20P; O–Chromatorex ODS; S–Sephadex LH-20; C–Semi-preparative reversed-phase Symmetry C-18 column; F–Fraction.	<b>67</b>
<b>Figure 3.1.3:</b> Compounds isolated from the leaves of <i>P. rotundifolia</i> and <i>P. praetermissa</i> .	<b>68</b>
<b>Figure 3.1.1.1:</b> Fragmentation pattern for prunasin 6'- <i>O</i> -gallate (1).	<b>72</b>
<b>Figure 3.1.1.2:</b> Full scan mass spectra ( $m/z$ ) with isotopic pattern for prunasin 6'- <i>O</i> -gallate (1).	<b>73</b>
<b>Figure 3.1.1.3:</b> Fragmentation pattern for prunasin 2',6'-di- <i>O</i> -gallate (2).	<b>75</b>
<b>Figure 3.1.1.4:</b> Full scan mass spectra ( $m/z$ ) with isotopic pattern for prunasin 2',6'-di- <i>O</i> -gallate (2).	<b>76</b>
<b>Figure 3.1.1.5:</b> Fragmentation pattern for prunasin 3',6'-di- <i>O</i> -gallate (3).	<b>78</b>
<b>Figure 3.1.1.6:</b> Full scan mass spectra ( $m/z$ ) with isotopic pattern for prunasin 3',6'-di- <i>O</i> -gallate (3).	<b>79</b>
<b>Figure 3.1.1.7:</b> Fragmentation pattern for prunasin 4',6'-di- <i>O</i> -gallate (4).	<b>81</b>
<b>Figure 3.1.1.8:</b> Full scan mass spectra ( $m/z$ ) with isotopic pattern for prunasin 4',6'-di- <i>O</i> -gallate (4).	<b>82</b>
<b>Figure 3.1.1.9:</b> Fragmentation pattern for prunasin 2',3',6'-tri- <i>O</i> -gallate (5).	<b>84</b>
<b>Figure 3.1.1.10:</b> Full scan mass spectra ( $m/z$ ) with isotopic pattern for prunasin 2',3',6'-tri- <i>O</i> -gallate (5).	<b>85</b>
<b>Figure 3.1.1.11:</b> Fragmentation pattern for prunasin 3',4',6'-tri- <i>O</i> -gallate (6).	<b>87</b>
<b>Figure 3.1.1.12:</b> Full scan mass spectra ( $m/z$ ) with isotopic pattern for prunasin 3',4',6'-tri- <i>O</i> -gallate (6).	<b>88</b>

<b>Figure 3.1.1.13:</b>	Fragmentation pattern for prunasin 2',3',4',6'-tetra- <i>O</i> -gallate (7).	<b>90</b>
<b>Figure 3.1.1.14:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for prunasin 2',3',4',6'-tetra- <i>O</i> -gallate (7).	<b>91</b>
<b>Figure 3.1.1.15:</b>	Fragmentation scheme for galloylated cyanogenic glucosides (1-7) observed by data dependent MS <sup>n</sup> with dynamic exclusion in negative mode.	<b>93</b>
<b>Figure 3.1.2.1:</b>	Fragmentation pattern for 6- <i>O</i> -galloyl-D-glucose (8).	<b>97</b>
<b>Figure 3.1.2.2:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 6- <i>O</i> -galloyl-D-glucose (8).	<b>98</b>
<b>Figure 3.1.2.3:</b>	Fragmentation pattern for 3,6-di- <i>O</i> -galloyl-D-glucose (9).	<b>100</b>
<b>Figure 3.1.2.4:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 3,6-di- <i>O</i> -galloyl-D-glucose (9).	<b>101</b>
<b>Figure 3.1.2.5:</b>	Fragmentation pattern for 1,2,3-tri- <i>O</i> -galloyl-β-D-glucose (10).	<b>103</b>
<b>Figure 3.1.2.6:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 1,2,3-tri- <i>O</i> -galloyl-β-D-glucose (10).	<b>104</b>
<b>Figure 3.1.2.7:</b>	Fragmentation pattern for 1,4,6-tri- <i>O</i> -galloyl-β-D-glucose (11).	<b>106</b>
<b>Figure 3.1.2.8:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 1,4,6-tri- <i>O</i> -galloyl-β-D-glucose (11).	<b>107</b>
<b>Figure 3.1.2.9:</b>	Fragmentation pattern for 3,4,6-tri- <i>O</i> -galloyl-D-glucose (12).	<b>109</b>
<b>Figure 3.1.2.10:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 3,4,6-tri- <i>O</i> -galloyl-D-glucose (12).	<b>110</b>
<b>Figure 3.1.2.11:</b>	Fragmentation pattern for 1,2,3,6-tetra- <i>O</i> -galloyl-β-D-glucose (13).	<b>112</b>
<b>Figure 3.1.2.12:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 1,2,3,6-tetra- <i>O</i> -galloyl-β-D-glucose (13).	<b>113</b>
<b>Figure 3.1.2.13:</b>	Fragmentation pattern for 1,2,3,4,6-penta- <i>O</i> -galloyl-β-D-glucose (14).	<b>115</b>
<b>Figure 3.1.2.14:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 1,2,3,4,6-penta- <i>O</i> -galloyl-β-D-glucose (14).	<b>116</b>

<b>Figure 3.1.2.15:</b>	Fragmentation pathway of gallotannins ( <b>8-14</b> ) observed by data dependent MS <sup>n</sup> with dynamic exclusion in negative mode. (The mass loss of 170, 152, 144, 60, 44 and 18 represented the removal of gallate, galloyl, hexosyl-H <sub>2</sub> O, 2CH <sub>2</sub> O, CO <sub>2</sub> and H <sub>2</sub> O respectively).	<b>118</b>
<b>Figure 3.1.3.1:</b>	Fragmentation pattern for 6- <i>O</i> -galloyl-2,3- <i>O</i> -( <i>S</i> )-hexahydroxydiphenoyl-D-glucose ( <b>15</b> ).	<b>123</b>
<b>Figure 3.1.3.2:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 6- <i>O</i> -galloyl-2,3- <i>O</i> -( <i>S</i> )-hexahydroxydiphenoyl-D-glucose ( <b>15</b> ).	<b>124</b>
<b>Figure 3.1.3.3:</b>	Fragmentation pattern for praecoxin B ( <b>16</b> ).	<b>126</b>
<b>Figure 3.1.3.4:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for praecoxin B ( <b>16</b> ).	<b>127</b>
<b>Figure 3.1.3.5:</b>	Fragmentation pattern for pterocarinin C ( <b>17</b> ).	<b>129</b>
<b>Figure 3.1.3.6:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for pterocarinin C ( <b>17</b> ).	<b>130</b>
<b>Figure 3.1.3.7:</b>	Fragmentation pattern for casuarinin ( <b>18</b> ).	<b>132</b>
<b>Figure 3.1.3.8:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for casuarinin ( <b>18</b> ).	<b>133</b>
<b>Figure 3.1.3.9:</b>	Fragmentation pattern of ellagitannins ( <b>15-17</b> ) in negative mode.	<b>135</b>
<b>Figure 3.1.4.1:</b>	Fragmentation pattern for 3'- <i>O</i> -methyl-3,4-methylenedioxyellagic acid 4'- <i>O</i> -β-D-glucopyranoside ( <b>19</b> ).	<b>139</b>
<b>Figure 3.1.4.2:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 3'- <i>O</i> -methyl-3,4-methylenedioxyellagic acid 4'- <i>O</i> -β-D-glucopyranoside ( <b>19</b> ).	<b>140</b>
<b>Figure 3.1.4.3:</b>	Fragmentation pattern for 3,3',4-tri- <i>O</i> -methylellagic acid 4'- <i>O</i> -β-D-glucopyranoside ( <b>20</b> ).	<b>142</b>
<b>Figure 3.1.4.4:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for 3,3',4-tri- <i>O</i> -methylellagic acid 4'- <i>O</i> -β-D-glucopyranoside ( <b>20</b> ).	<b>143</b>
<b>Figure 3.1.5.1:</b>	Fragmentation pattern for gallic acid ( <b>21</b> ).	<b>145</b>
<b>Figure 3.1.5.2:</b>	Full scan mass spectra ( <i>m/z</i> ) with isotopic pattern for gallic acid ( <b>21</b> ).	<b>146</b>

<b>Figure 3.1.5.3:</b>	Fragmentation pattern for gallic acid methyl ester ( <b>22</b> ).	<b>147</b>
<b>Figure 3.1.5.4:</b>	Full scan mass spectra ( $m/z$ ) with isotopic pattern for gallic acid methyl ester ( <b>22</b> ).	<b>148</b>
<b>Figure 3.2.1.1:</b>	<b>a)</b> ID-IR spectra ( $4000\text{-}450\text{ cm}^{-1}$ ) and <b>b)</b> Expanded spectra ( $1900\text{-}850\text{ cm}^{-1}$ ) for <i>P. praetermissa</i> and <i>P. rotundifolia</i> collected from different localities.	<b>151</b>
<b>Figure 3.2.2.1:</b>	Representative second derivative spectra for <i>P. rotundifolia</i> and <i>P. praetermissa</i> collected from different localities in the region of <b>a)</b> $1800\text{-}1600\text{ cm}^{-1}$ and <b>b)</b> $1300\text{-}800\text{ cm}^{-1}$ .	<b>155</b>
<b>Figure 3.2.3.1:</b>	PCA scores plot of <i>P. rotundifolia</i> collected from different localities.	<b>158</b>
<b>Figure 3.2.3.2:</b>	PCA scores plot of <i>P. praetermissa</i> collected from different localities.	<b>159</b>
<b>Figure 3.2.3.3:</b>	PCA scores plot of <i>P. rotundifolia</i> and <i>P. praetermissa</i> collected from each location.	<b>160</b>
<b>Figure 3.2.3.4:</b>	PCA scores plot of <b>a)</b> <i>P. rotundifolia</i> (red) and <i>P. praetermissa</i> (blue) from <b>b)</b> different localities ( <b>A:</b> <i>P. rotundifolia</i> from Pasoh Forest Reserve, Negeri Sembilan; <b>B:</b> <i>P. rotundifolia</i> from Sungai Batang, Labis Forest Reserve, Johor; <b>C:</b> <i>P. rotundifolia</i> from Takar Melor, Labis Forest Reserve, Johor; <b>D:</b> <i>P. praetermissa</i> from Bukit Lagong, Selangor; <b>E:</b> <i>P. praetermissa</i> from Ampang Forest Reserve, Selangor; <b>F:</b> <i>P. praetermissa</i> from Pasoh Forest Reserve, Negeri Sembilan).	<b>161</b>
<b>Figure 3.2.4.1:</b>	Synchronous 2D-IR contour plot ( <b>a</b> ), synchronous 2D-IR auto peaks ( <b>b</b> ) and synchronous 2D-IR mesh plot ( <b>c</b> ) in the range of $1500\text{-}1000\text{ cm}^{-1}$ for <i>P. rotundifolia</i> from Takar Melor, Labis Forest Reserve, Johor ( <b>PRLTM</b> ), Sungai Batang, Labis Forest Reserve, Johor ( <b>PRLSB</b> ) Pasoh Forest Reserve ( <b>PRLPA</b> ), and <i>P. praetermissa</i> from Pasoh Forest Reserve ( <b>PPLPA</b> ), Ampang Forest Reserve ( <b>PPLAM</b> ), Bukit Lagong ( <b>PPLLA</b> ).	<b>165</b>

<b>Figure 3.2.5.1:</b>	a) 1D-IR spectra (4000-450 cm <sup>-1</sup> ) and b) second derivative spectra (1300-800 cm <sup>-1</sup> ) for the mixture of <i>P. rotundifolia</i> ( <b>PRL</b> ) from Pasoh Forest Reserve and <i>P. praetermissa</i> ( <b>PPL</b> ) from Bukit Lagong.	<b>168</b>
<b>Figure 3.2.5.2:</b>	PCA scores plot obtained from 2000-450 cm <sup>-1</sup> IR spectra. The percentage indicated the adulteration of <i>P. praetermissa</i> ( <b>PPL</b> ) from Bukit Lagong in <i>P. rotundifolia</i> ( <b>PRL</b> ) from Pasoh Forest Reserve.	<b>169</b>
<b>Figure 3.2.5.3:</b>	Synchronous 2D-IR contour plot ( <b>A</b> ), synchronous 2D-IR auto peaks ( <b>B</b> ) and synchronous 2D-IR mesh plot ( <b>C</b> ) in the range of 1500-1000 cm <sup>-1</sup> for the mixture of <i>P. rotundifolia</i> from Pasoh Forest Reserve ( <b>PRLPA</b> ) and <i>P. praetermissa</i> from Bukit Lagong ( <b>PPLLA</b> ).	<b>170</b>
<b>Figure 3.3.1.1:</b>	HPLC profiles from the leaves of <i>P. rotundifolia</i> and <i>P. praetermissa</i> from different localities.	<b>172</b>
<b>Figure 3.3.2.1:</b>	HPLC chromatogram of ten individuals for <i>P. rotundifolia</i> and <i>P. praetermissa</i> from different localities after alignment.	<b>176</b>
<b>Figure 3.3.2.2:</b>	PCA scores plot of <i>P. rotundifolia</i> chromatographic profiles from different localities.	<b>177</b>
<b>Figure 3.3.2.3:</b>	PCA scores plot of <i>P. praetermissa</i> chromatographic profiles from different localities.	<b>178</b>
<b>Figure 3.3.2.4:</b>	PCA scores plot of <i>P. rotundifolia</i> and <i>P. praetermissa</i> chromatographic profiles from each location.	<b>179</b>
<b>Figure 3.4.1.1:</b>	LC-MS chromatographic profile of a) <i>P. rotundifolia</i> (Pasoh Forest Reserve) and b) expanded chromatogram.	<b>182</b>
<b>Figure 3.4.1.2:</b>	LC-MS chromatographic profile of a) <i>P. rotundifolia</i> (Sungai Batang, Labis Forest Reserve) and b) expanded chromatogram.	<b>183</b>
<b>Figure 3.4.1.3:</b>	LC-MS chromatographic profile of a) <i>P. rotundifolia</i> (Takar Melor, Labis Forest Reserve) and b) expanded chromatogram.	<b>184</b>
<b>Figure 3.4.1.4:</b>	LC-MS chromatographic profile of a) <i>P. praetermissa</i> (Bukit Lagong) and b) expanded chromatogram.	<b>185</b>

<b>Figure 3.4.1.5:</b>	LC-MS chromatographic profile of <b>a)</b> <i>P. praetermissa</i> (Ampang Forest Reserve) and <b>b)</b> expanded chromatogram.	<b>186</b>
<b>Figure 3.4.1.6:</b>	LC-MS chromatographic profile of <b>a)</b> <i>P. praetermissa</i> (Pasoh Forest Reserve) and <b>b)</b> expanded chromatogram.	<b>187</b>
<b>Figure 3.4.1.7:</b>	LC-MS chromatographic profiles of <i>P. rotundifolia</i> and <i>P. praetermissa</i> from different localities.	<b>188</b>
<b>Figure 3.5.1:</b>	Neuroprotective activity of galloylated cyanogenic glucosides ( <b>1-7</b> ) in bar graph. * <i>P</i> < 0.01 was considered to be significant.	<b>195</b>
<b>Figure 3.5.2:</b>	Neuroprotective activity of gallotannins ( <b>8-14</b> ) in bar graph. * <i>P</i> < 0.01 was considered to be significant.	<b>196</b>
<b>Figure 3.5.3:</b>	Neuroprotective activity of ellagitannins ( <b>15-18</b> ), ellagic acid derivatives ( <b>19-20</b> ), gallic acid ( <b>21</b> ) and gallic acid methyl ester ( <b>22</b> ). * <i>P</i> < 0.01 was considered to be significant.	<b>197</b>
<b>Figure 3.6.1.1:</b>	<i>In vitro</i> cytotoxicity on CaSki cell line of galloylated cyanogenic glucosides ( <b>1-7</b> ). * <i>P</i> < 0.05 was considered to be significant.	<b>199</b>
<b>Figure 3.6.1.2:</b>	<i>In vitro</i> cytotoxicity on CaSki cell line of gallotannins ( <b>8-14</b> ). * <i>P</i> < 0.05 was considered to be significant.	<b>200</b>
<b>Figure 3.6.1.3:</b>	<i>In vitro</i> cytotoxicity on CaSki cell line of ellagitannins ( <b>15-18</b> ), ellagic acid derivatives ( <b>19-20</b> ) and aromatic compounds ( <b>21-22</b> ). * <i>P</i> < 0.05 was considered to be significant.	<b>201</b>
<b>Figure 3.6.2.1:</b>	<i>In vitro</i> cytotoxicity on HCT 116 cell line of galloylated cyanogenic glucosides ( <b>1-7</b> ). * <i>P</i> < 0.05 was considered to be significant.	<b>203</b>
<b>Figure 3.6.2.2:</b>	<i>In vitro</i> cytotoxicity on HCT 116 cell line of gallotannins ( <b>8-14</b> ). * <i>P</i> < 0.05 was considered to be significant.	<b>204</b>
<b>Figure 3.6.2.3:</b>	<i>In vitro</i> cytotoxicity on HCT 116 cell line of ellagitannins ( <b>15-18</b> ), ellagic acid derivatives ( <b>19-20</b> ) and aromatic compounds ( <b>21-22</b> ). * <i>P</i> < 0.05 was considered to be significant.	<b>205</b>

- Figure 3.6.3.1:** *In vitro* cytotoxicity on MCF 7 cell line of galloylated cyanogenic glucosides (1-7). \* $P < 0.05$  was considered to be significant. 207
- Figure 3.6.3.2:** *In vitro* cytotoxicity on MCF 7 cell line of gallotannins (8-14). \* $P < 0.05$  was considered to be significant. 208
- Figure 3.6.3.3:** *In vitro* cytotoxicity on MCF 7 cell line of ellagitannins (15-18), ellagic acid derivatives (19-20) and aromatic compounds (21-22). \* $P < 0.05$  was considered to be significant. 209