

REFERENCES

- Adli, D. S. H., Stusse, S. L., and Cruce, W. L. R. (1999). Immunohistochemistry and spinal projections of the reticular formation in the northern leopard frog, *Rana pipiens*. *Journal of Comparative Neurology*, 404: 387-407.
- Adli, D. S. H., and Cruce, W. L. R. (1995a). Immunohistochemical localization of serotonin in the brain stem of *Rana pipiens* (Amphibia). *Malaysian Journal of Science*, 16A: 69-75.
- Adli, D. S. H., and Cruce, W. L. R. (1995b). Neural circuitry revealed by immunohistochemistry. *Malaysian Journal of Medical Laboratory Sciences*, 12(1): 15-19.
- Adli, D. S. H., Rosenthal, B. M., Yuen, G. L., Ho, R. H., and Cruce, W. L. R. (1988). Immunohistochemical localization of substance P, somatostatin, enkephalin, and serotonin in the spinal cord of the Northern Leopard frog, *Rana pipiens*. *The Journal of Comparative Neurology*, 275:106-116.
- Aronin, N, Difiglia, M., and Leeman, S. E. (1983). Substance P. In Krieger, D. T., Brownstein, M. J. and Martin, J. B. (Eds.), *Brain peptides* (pp. 783-801). New York: Wiley Interscience.
- Atweh, S. F., and Kuhar, M. J. (1977). Autoradiographic localization of opiate receptors in rat brain. I. Spinal cord and lower medulla. *Brain Research*, 124(1): 53-67.
- Azlinda Abu Bakar. (1996). *Perbandingan korda spina antara dua spesies haiwan, Bufo melanostictus dan Periophthalmodon schlosseri*. Thesis of the Bachelor of Science Degree. Zoology Department, Faculty of Science, University of Malaya, Kuala Lumpur.
- Berry, P. Y. (1975). *The amphibian fauna of Peninsular Malaysia*. Kuala Lumpur: Tropical Press.
- Bertheau, P., Cazals-Hatem, D., Meignin, V., de Roquancourt, A., Vérola, O., Lesourd, A., et al. (1998). Variability of immunohistochemical reactivity on stored paraffin slides. *Journal of Clinical Pathology*, 51(5): 370-374
- Brinkman, R. and Martin, A. H. (1973). A cytoarchitectonic study of the spinal cord of the domestic fowl *Gallus gallus domesticus*. I. brachial region. *Brain Research*, 56: 43-62

Boenisch, T. (Ed.). (2001). *Dako Handbook: Immunohistochemical Staining Methods* (3rd ed.). California: Dako Corporation.

Buchwalow, I. B., and Böcker, W. (2010). *Immunohistochemistry: basics and methods*. Berlin: Springer Verlag.

Caamaño-Tubío, R. I., Pérez, J., Ferreiro, S., and Aldegunde, M. (2007). Peripheral serotonin dynamics in the rainbow trout (*Oncorhynchus mykiss*). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 145(2): 245-255

Cakir, Y., and Strauch, S. M. (2005). Tricaine (MS-222) is a safe anesthetic compound compared to benzocaine and pentobarital to induce anesthesia in leopard frogs (*Rana pipiens*). *Pharmacological Reports*, 57: 467-474.

Cameron, A. A., Plenderleith, M. B. and Snow, P. J. (1990). Organization of the spinal cord in four species of elasmobranch fishes: cytoarchitecture and distribution of serotonin and selected neuropeptides. *The Journal of Comparative Neurology*, 297: 201-218

Carlin, K. P., Dai, Y., and Jordan, L. M. (2006). Cholinergic and serotonergic excitation of ascending commissural neurons in the thoraco-lumbar spinal cord of the neonatal mouse. *Journal of Neurophysiology*, 95: 1278-1284.

Cesselin, F., Bourgoin, S., Clot, A. M., Hamon, M., and Le Bars, D. (1989). Segmental release of Met-enkephalin-like material from the spinal cord of rats, elicited by noxious thermal stimuli. *Brain Research*, 484(1-2): 71-77.

Chan Palay, V., Jonsson, G., and Palay, S. L. (1978). Serotonin and substance P coexist in neurons of the rat's central nervous system. *Proceedings of the National Academy of Sciences USA*, 75(3): 1582-1586.

Chan-Palay, V., and Palay, S. L. (1977). Ultrastructural identification of substance P cells and their processes in rat sensory ganglia and their terminals in the spinal cord by immunocytochemistry. *Proceedings of the National Academy of Sciences USA*, 74(9): 4050-4054.

Conrath-Verrier, M., Dietl, M., Arluison, M., Cesselin, F., Bourgoin, S., and Hamon, M. (1983). Localization of met-enkephalin-like immunoreactivity within pain-related nuclei of cervical spinal cord, brainstem and midbrain in the cat. *Brain Research Bulletin*, 11(5): 587-604

Crisp, T., Stafinsky, J. L., Spanos, L. J., Uram, M., Perni, V. C., and Donepudi, H. B. (1991). Analgesic effects of serotonin and receptor-selective serotonin agonists in the rat spinal cord. *General Pharmacology: The Vascular System*, 22(2): 247-251

Crossman, A. R., and Neary, D. (2005). *Neuroanatomy: An illustrated colour text* (3rd ed.). Manchester: Elsevier Churchill Livingstone.

Cruce, W. L. R. (1974). A suprasinal monosynaptic input to hindlimb motoneurons in the lumbar spinal cord of the frog, *Rana catesbeiana*. *Journal of Neurophysiology*, 37(4): 691-704

Cruce, W. L. R. (1979). Spinal cord in lizards. In Gans, C., Northcutt, R. G. and Ulinski P. S. (Eds.). *Biology of the Reptilia* (pp. 111-132). New York: Academic Press.

Cullen, K. E. (2009). *Encyclopedia of Life Science* (Vol. 1). New York: Infobase Publishing.

Dafny, N. (1997). *Anatomy of the spinal cord*. Retrieved October 19, 2011 from The University of Texas Health Science Center website: <http://neuroscience.uth.tmc.edu/s2/chapter03.html>

Davis, B. M., and Cabot, J. B. (1984). Substance P-containing pathways to avian sympathetic preganglionic neurons: evidence for major spinal-spinal circuitry. *The Journal of Neuroscience*, 4(8): 2145-2159.

De Carlos J.A., and Borrel J. (2007). A historical reflection of the contributions of Cajal and Golgi to the foundations of neuroscience. *Brain Research Reviews*, 55: 8-16.

De Falco, M., Laforgia, V., Valiante, S., Virgilio, F., Varano, L., and De Luca, A. (2002). Different patterns of expression of five neuropeptides in the adrenal gland and kidney of two species of frog. *The Histochemical Journal*, 34: 221-26.

De Felipe, C., Herrero, J. F., O'Brien, J. A., Palmer, J. A., Doyle, C. A., Smith, A. J. H., et al. (1998). Altered nociception, analgesia and aggression in mice lacking the receptor for substance P. *Nature*, 392: 394-397.

De Koninck, Y., Ribeiro-da-Silva, A., Henry, J. L., and Cuello, A. C. (1992). Spinal neurons exhibiting a specific nociceptive response receive abundant substance P-containing synaptic contacts. *Proceedings of the National Academy of Sciences USA*, 89(11): 5073-5077.

Droge, M. H., and Leonard, R. B. (1981). Organization of spinal motor nuclei in the stingray, *Dasyatis sabina*. *Brain Research*, 276(2): 201-211.

Dubé, D., and Pelletier, G. (1979). Effect of colchicine on the immunohistochemical localization of somatostatin in the rat brain: light and electron microscopic studies. *The Journal of Histochemistry and Cytochemistry*, 27(12): 1577-1581.

Du, F., and Dubois, P. M. (1988). Development and distribution of enkephalin-immunoreactive elements in the chicken spinal cord. *Neuroscience*, 27(1): 251-266.

Ebbeson, S. O. E. (1976). Spinal cord. In Llinas, R. and Precht, W. (Eds.). *Frog neurobiology: a handbook* (pp. 679-706). New York: Springer-Verlag.

Ecology Asia: Field Frog. (2008). Retrieved November 14, 2008 from http://ecologyasia.com/verts/amphibians/field_frog.htm

Elde, R. (1985). Immunocytochemistry. In Krieger, D. T., Brownstein, M. J. and Martin, J. B. (Eds.). *Brain peptides* (pp. 485-493). New York: Wiley Interscience.

Fernández, A., Radmilovich, M. and Trujillo-Cenóz, O. (1992). Neuronal types in the spinal dorsal gray of the turtle *Chrysemys d'orbigny*: a Golgi study. *Cell Tissue Research*, 272: 355-368.

FitzGerald, M. J. T., Gruener, G., and Mtui, E. (2007). *Clinical neuroanatomy and neuroscience* (5th ed.). Philadelphia: Saunders Elsevier.

FitzGerald, M. J. T. (1985). *Neuroanatomy basic & applied*. England: Baillière Tindall.

Fürst, S. (1999). Transmitters involved in antinociception in the spinal cord. *Brain Research Bulletin*, 48(2): 129-141.

Galhardo, V. and Lima, D. (1999). Structural characterization of marginal (lamina I) spinal cord neurons in the cat: a Golgi study. *Journal of Comparative Neurology*, 414: 315-333.

Gamse, R., Lackner, D., Gamse, G., and Leeman, S. E. (1981). Effect of capsaicin pretreatment on capsaicin-evoked release of immunoreactive somatostatin and substance P from primary sensory neurons. *Naunyn-Schmiedeberg's Archives of Pharmacology*, 316(1): 38-41

Gamse, R., Molnar, A., and Lembeck, F. (1979). Substance P release from spinal cord slices by capsaicin. *Life Sciences*, 25(7): 629-636.

Glazer, E. J., and Basbaum, A. I. (1981). Immunohistochemical localization of leucine-enkephalin in the spinal cord of the cat: Enkephalin-containing marginal neurons and pain modulation. *The Journal of Comparative Neurology*, 196(3): 377-389.

Glazer, E. J., and Basbaum, A. I. (1983). Opioid neurons and pain modulation: an ultrastructural analysis of enkephalin in cat superficial dorsal horn. *Neuroscience*, 10(2): 357-76.

Goodman, R. R., Fricker, L. D., and Snyder, S. H. (1983). Enkephalins. In Krieger, D. T., Brownstein, M. J. and Martin, J. B. (Eds.), *Brain peptides* (pp. 783-801). New York: Wiley Interscience.

Grant, G., and Koerber, H. R. (2004). Spinal cord cytoarchitecture. In Paxinos, G. (Ed.), *The rat spinal cord* (3rd ed.) (pp. 121-128). USA: Elsevier.

Gravenhorst, J. L. C. (1829). Deliciae Musei Zoologici Vratislaviensis. Fasciculus Primus, Continens Chelonios et Batrachiae. Sumptibus Leopoldi Vossii. Lipsiae.

Gray, E. G. (1957). The spindle and extrafusal innervations of a frog muscle. *Proc. R. Soc. Lond. B. Biol. Sci.*, 146: 416-430.

Guedes, R. P., Marchi, M. I., Viola, G. G., Xavier, L. L., Achaval, M., and Partata, W. A. (2004). Somatostatin-, calcitonin gene-related peptide, and γ -aminobutyric acid-like immunoreactivity in the frog lumbosacral spinal cord: distribution and effects of sciatic nerve transaction. *Comparative Biochemistry and Physiology*, 138: 19-28.

Gurr, E. (1956). *A practical manual of medical & biological staining techniques*. London: Leonard Hill Limited.

Hadjiconstantinou, M., Panula, P., Lackovic, Z., and Neff, N. H. (1984). Spinal cord serotonin: a biochemical and immunohistochemical study following transection. *Brain Research*, 322: 245-254.

Hancock, M. B. (1982). A serotonin-immunoreactive fiber system in the dorsal columns of the spinal cord. *Neuroscience Letters*, 31(3): 247-252.

Halpern, M. (1972). Some connections of the telencephalon of the frog, *Rana pipiens*. *Brain, Behavior and Evolution*, 6: 42-68.

Heise, C., and Kayalioglu, G. (2009). Cytoarchitecture of the spinal cord. In Watson, C., Paxinos, G. and Kayalioglu, G. (Eds.), *The spinal cord. A Christopher and Dana Reeve Foundation Text and Atlas* (pp. 64-93). United Kingdom: Academic Press.

Hendelman, W. J. (2000). *Atlas of functional neuroanatomy*. Florida: CRC Press.

Ho, R. H., and LaValley, A. L. (1984). 5-Hydroxytryptamine in the spinal cord of the domestic fowl. *Brain Research Bulletin*, 13: 427-431.

Hochman, S. (2007). Spinal cord. *Current Biology*, 17(2): 950-955.

Holland, M. S., Mackenzie, C. D., Bull, R. W., and Silva, R. F. (1996). A comparative study of histological conditions suitable for both immunofluorescence and in situ hybridization in the detection of Herpesvirus and its antigens in chicken tissues. *Journal of Histochemistry and Cytochemistry*, 44(3):259-65.

Honda, C. N., and Lee, C. L. (1985). Immunohistochemistry of synaptic input and functional characterizations of neurons near the spinal central canal. *Brain Research*, 343(1): 120-128.

Hökfelt, T., Kellerth, J., Nilsson, G., and Pernow, B. (2003). Experimental immunohistochemical studies on the localization and distribution of substance P in cat primary sensory neurons. *Brain Research*, 100(2): 235-252.

Hökfelt, T., Ljungdahl, A., Terenius, L., Elde, R., and Nilsson, G. (1977). Immunohistochemical analysis of peptide pathways possibly related to pain and analgesia: Enkephalin and substance P. *Proceedings of the National Academy of Sciences USA*, 74(7): 3081-3085.

Hökfelt, T., Terenius, L., Kuypers, H. G. J. M., and Dann, O. (1979). Evidence for enkephalin immunoreactive neurons in the medulla oblongata projecting to the spinal cord. *Neuroscience Letters*, 14(1): 55-60.

Huang, J. H., and Weiss, M. L. (1999). Characterization of the central cell groups regulating the kidney in the rat. *Brain Research*, 845(1): 77-91.

Inderbir S. (2006). *Textbook of human neuroanatomy* (7th ed.). New Delhi: Jaypee Brothers Medical Publisher (P) Ltd.

Inger, R. F., and Stuebing, R. B. (2005). *A field guide to the frogs of Borneo*. Kota Kinabalu: Natural History Publications.

Inman, D. M., and Steward, O. (2003). Ascending sensory, but not other long-tract axons, regenerate into the connective tissue matrix that forms at the site of a spinal cord injury in mice. *The Journal of Comparative Neurology*, 462(4): 431–449.

Iritani, S., Tohgi, M., Arai, T., and Ikeda, K. (2006). Immunohistochemical study of the serotonergic neuronal system in an animal model of mood disorder. *Experimental Neurology*. 201, 60-65.

Johansson, O., and Hökfelt, T. (1980). Thyrotropin releasing hormone, somatostatin, and enkephalin: Distribution studies using immunohistochemical techniques. *The Journal of Histochemistry and Cytochemistry*, 28(4): 364-366.

Kandel, E. R., and Schwartz, J. H. (1985). *Principles of neural science* (2nd ed.). New York: Elvevier.

Kanner, B. I., and Schuldiner, S. (1987). Mechanism of Transport and Storage of Neurotransmitter. *Critical reviews in Biochemistry and Molecular Biology*, 22(1):1-38.

Kantner, R. M., Kirby, M. L., and Goldstein, B. D. (1985). Increase in substance P in the dorsal horn during a chemogenic nociceptive stimulus. *Brain Research*, 338(1): 196-199.

Kirmani, F., Ghaus, F., and Ullah, M. (2011). Neuronal arrangement in cervical enlargement of spinal cord of rabbit. *Current Neurobiology*, 2 (1): 53-56

Kojima, M., Takeuchi, Y., Goto, M., and Sano, Y. (1983) Immunohistochemical study on the localization of serotonin fibers and terminals in the spinal cord of the monkey. *Cell Tissue Research*, 229: 23-36.

Kuhlenbeck, H. (1975). *The central nervous system of vertebrates: a general survey of its comparative anatomy with an introduction to the pertinent fundamental biologic and logical concepts*. New York: Karger Publisher.

- Kuraishi, Y., Hirota, N., Sato, Y., Hino, Y., Satoh, M., and Takagi, H. (1985). Evidence that substance P and somatostatin transmit separate information related to pain in the spinal dorsal horn. *Brain Research*, 325: 294-298.
- Kusuma, A., Ten Donkelaar, H. J., and Nieuwenhuys, R. (1979). Intrinsic organization of the spinal cord. In Gans, C., Northcutt, R. G. and Ulinski P. S. (Eds.). *Biology of the Reptilia* (pp. 59-107). New York: Academic Press.
- LaMotte, C. C. (1988). Lamina X of primate spinal cord: distribution of five neuropeptides and serotonin. *Neuroscience*, 25(2): 639-658.
- Lanerolle, N. C., and LaMotte, C. C. (1982). The human spinal cord: Substance P and methionine-enkephalin immunoreactivity. *The Journal of Neuroscience*, 2(10): 1369-1386.
- Lautenbacher, S., and Fillingim, R. B (Eds.). (2004). *Pathophysiology of pain perception*. New York: Plenum Publisher.
- LaValley, A. (1980). *Distribution of substance P (SP), somatostatin (SOM) and met-enkephalin immunoreactivities in the spinal cord of the domestic fowl, Gallus domesticus*. Thesis of Degree Master of Science, Ohio State University, the United States.
- LaValley, A. L., and Ho, R. H. (1983). Substance P, somatostatin and methionine enkephalin-immunoreactive elements in the spinal cord of the domestic fowl. *The Journal of Comparative Neurology*, 213(4): 406-413.
- Lembeck, F., Donnerer, J., and Colpaert, F.C. (1981). Increase of substance P in primary afferent nerves during chronic pain. *Neuropeptides*, 1(3):175-180.
- Leonard, R. B., and Cohen D. H. (1975). A cytoarchitectonic analysis of the spinal cord of the pigeon (*Columba livia*). *The Journal of Comparative Neurology*, 163(2): 159-180.
- Light, A. R., and Metz, C. B. (1978). The morphology of the spinal cord efferent and afferent neurons contributing to the ventral roots of the cat. *The Journal of Comparative Neurology*, 179(3): 501-515.
- Lillie, R. D. (1954). *Histopathologic technic and practical histochemistry*. Toronto: McGraw-Hill Book Company.

Lima, D., Avelina, A., and Coimbra, A. (1993). Morphological characterization of marginal (lamina I) neurons immunoreactive for substance P, enkephalin, dynorphin and gamma-aminobutyric acid in rat spinal cord. *Journal of Chemical Neuroanatomy*, 6: 43-52.

Llinas, R., and Precht, W. (Eds.). (1976) *Frog neurobiology: a handbook*. New York: Springer-Verlag.

Lorez, H. P., and Kemali, M. (1981). Substance P-, met-enkephalin- and somatostatin-like immunoreactivity distribution in the frog spinal cord. *Neuroscience Letters*, 26: 110-124.

Ma, W., Ribeiro-da-Silva, A., De Koninck, Y., Radhakrishnan, V., Cuello, A. C., and Henry J. L. (1997). Substance P and enkephalin immunoreactivities in axonal boutons presynaptic to physiologically identified dorsal horn neurons. An ultrastructural multiple-labelling study in the cat. *Neuroscience*, 77(3): 793-811.

Maccio-Hage, I. (2005). *Pain in fish*. Retrieved May 12, 2009, from <http://www.fair-fish.ch>

Machin, K. L. (1999). Amphibian pain and analgesia. *Journal of Zoo and Wildlife Medicine*, 30, 2-20.

Marshall, A. J. (Ed.). (1961), *Biology and comparative physiology of birds* (Vol. 2). New York: Academic Press.

McClung, J. R., and Castro, A. J. (1978) Rexed's laminar scheme as it applies to the rat cervical spinal cord. *Experimental Neurology* 58: 145–148.

McDiarmid, R. W., and Altig, R. (1999). *Tadpoles: the biology of anuran larvae*. United States of America: University of Chicago Press.

Mensah, P. L., and Thompson, R. F. (1978). Descending fibres of the lateral funiculus of the amphibian spinal cord: their course and terminal distribution. *Journal of Anatomy*, 125(1): 1-9.

Messing, R. B., and Lytle, L. D. (1977). Serotonin-containing neurons: their possible role in pain and analgesia. *Pain*, 4: 1-21

Micevych, P. E., Coquelin, A., and Arnold, A. P. (1986). Immunohistochemical distribution of substance P, serotonin, and methionine enkephalin in sexually

dimorphic nuclei of the rat lumbar spinal cord. *The Journal of Comparative Neurology*, 248(2): 235–244.

Miller, R. T. (2001). *Technical immunohistochemistry: achieving reliability and reproducibility of immunostains*. Retrieved June 24, 2011 from IHCWorld website: http://www.ihcworld.com/_books/books.htm.

Miller, R. J., and Pickel, V. M. (1980). The distribution and functions of the enkephalin. *The Journal of Histochemistry and Cytochemistry*, 28(8): 903-917.

Miller, K. E., and Seybold, V. S. (1987). Ultrastructure of the central gray region (lamina X) in cat spinal cord. *Neuroscience*, 22(3): 1057-1066.

Molander, C., Xu, Q., and Grant, G. (1984). The cytoarchitectonic organization of the spinal cord in the rat. I. The lower thoracic and lumbosacral cord. *The Journal of Comparative Neurology*, 230(1): 133–141.

Naftchi, N. E., Abrahams, S. J., St. Paul, H. M., Lowman, E. W., and Scholosser, W. (1978). Localization and changes of substance P in spinal cord of paraplegic cats. *Brain Research*, 153(3): 507-513.

Newton, B. W., Maley, B., and Hamill, R. W. (1986). Immunohistochemical demonstration of serotonin neurons in autonomic regions of the rat spinal cord. *Brain Research*, 376: 155-163.

Nishikawa, K. C., Roth, G. and Dicke, U. (1991). Motor neurons and motor columns of the anterior spinal cord of salamanders: post-hatching development and phylogenetic distribution. *Brain, Behavior and Evolution*, 37: 368-382.

Noble, G. K. (1954). *The biology of amphibian*. New York: Dover Publications.

Oksche, A., and Ueck, M. (1976). The Nervous System. In Loft, B. (Eds.). *Physiology of the amphibian* (Vol. 3) (pp. 314-404). London: Academic Press.

Ottoson, D. (1983). *Physiology of the nervous system*. New York: Oxford University Press.

Oyadeyi, A. S., Ajao, F. O. Afolabi, A. O., Udoh, U. S., and Azeez, O. M. (2007). The formalin test in African Toad (*Bufo regularis*) – a novel pain model in amphibians. *American-Eurasian Journal of Scientific Research*, 2 (1): 24-28.

- Oyama, T., Ueda, M., Kuraishi, Y., Akaike, A., and Satoh, M. (1996). Dual effect of serotonin on formalin-induced nociception in the rat spinal cord. *Neuroscience Research*, 25(2): 129-35.
- Palkovitz, M. (1985). Neuroanatomical techniques. In Krieger, D. T., Brownstein, M. J. and Martin, J. B. (Eds.). *Brain peptides* (pp. 496-540). New York: Wiley Interscience.
- Partata, W. A., Cerveira, J. F., Xavier, L. L., Viola, G. G., and Achaval, M. (2002). Sciatic nerve transaction decrease substance P immunoreactivity in the lumbosacral spinal cord of the frog (*Rana catesbeiana*). *Comparative Biochemistry and Physiology*, 131: 80-814.
- Partata, W. A., Krepsky, A. M., Xavier, L. L., Marques, M., and Achaval, M. (2003). Substance P immunoreactivity in the lumbar spinal cord of the turtle *Trachemys dorbignyi* following peripheral nerve injury. *Brazilian Journal of Medical and Biological Research*, 36(4): 515-520.
- Pearce, J. M. S. (2008). The development of spinal cord anatomy. *European Neurology*, 59: 286-291.
- Peters, A. (2007). Golgi, Cajal, and the fine structure of the nervous system. *Brain Research Reviews*, 55(2): 256-263.
- Prescott, S. A., and De Koninck, Y. (2002). Four cell types with distinctive membrane properties and morphologies in lamina I of the spinal dorsal horn of the rat. *The Journal of Physiology*, 539: 817-836.
- Purkiss, J., Welch, M., Doward, S., and Foster, K. (2000). Capsaicin-stimulated release of substance P from cultured dorsal root ganglion neurons: involvement of two distinct mechanisms. *Biochemical Pharmacology*, 59(11): 1403-1406.
- Ralston, H. J. III. (1979). The fine structure of laminae I, II and III of the macaque spinal cord. *The Journal of Comparative Neurology*, 184 (4): 619–641.
- Reiner, A. (1987). The distribution of proenkephalin-derived peptides in the central nervous system of turtles. *The Journal of Comparative Neurology*, 259: 65-91.
- Rexed B. (1952). The cytoarchitectonic organization of the spinal cord in the cat. *The Journal of Comparative Neurology*, 96 (3): 414–95.

- Ribeiro-da-Silva, A., Pioro, E. P., and Cuello, A. C. (1991). Substance P- and enkephalin-like immunoreactivities are colocalized in certain neurons of the substantia gelatinosa of the rat spinal cord: an ultrastructural double-labeling study. *Journal of Neuroscience*, 11(4): 1068-1080.
- Robinson, G., Ellis, I. O., and MacLennan, K. A. (1990). Immunocytochemistry. In Bancroft, J. D. and Tumer, D. R. (Eds.), *Theory and practice of histological techniques* (3rd ed.) (pp. 413-433). United Kingdom: Churchill Livingstone.
- Rohaya Mamat. (1996). *Perkembangan system saraf pusat katak tempatan, Polypedates leucomystax (Boie) dari peringkat berudu ke peringkat dewasa*. Thesis of the Bachelor of Science Degree. Zoology Department, Faculty of Science, University of Malaya, Kuala Lumpur.
- Rolls, G. (2012). *Fixation and Fixatives (1) – The Process of Fixation and the Nature of Fixatives*. Retrieved May 21, 2012 from Leica Microsystems website: <http://www.leica-microsystems.com/pathologyleaders/fixation-and-fixatives-1-the-process-of-fixation-and-the-nature-of-fixatives/>
- Sahelian, R. (2005). *Neurotransmitters*. Retrieved November 11, 2008, from <http://www.raysahelian.com/neurotransmitter.html>
- Sakamoto, H. and Atsumi, S. (1989). Substance P-immunoreactive elements in laminae I and II of the chicken spinal cord: a light- and electron-microscopic study. *Neuroscience Research*, 7: 103-116.
- Santafé, M. M., Garcia N., Lanuza, M. A., Tomàs, M., Besalduch, N., Sabaté, M., and Tomàs, J. (2007). Histology in neuroscience. From the symposia: Homage of the histology chairs to Santiago Ramón y Cajal, 18–19 December, 2006, Albacete, Spain. *Journal of Molecular Histology*. 39(3): 1567-2387.
- Sarnat, H. B., and Netsky, M. G. (1981). *Evolution of the nervous system* (2nd ed.). New York: Oxford University Press.
- Sasaki, H. (1977). Cytoarchitectonic analysis of the bullfrog (*Rana catesbeiana*) spinal cord by means of electron microscopy with special reference to distribution of microneurons. *The Journal of Comparative Neurology*, 176: 101-120.
- Schaeffer, D. O. and Waters, R. M. (1996). Neuroanatomy and neurological diseases of reptiles. *Seminars in Avian and Exotic Pet Medicine*, 5(3): 165-171.

- Senba, E., Yanaihara, C., Yanaihara, N., and Tohyama, M. (1988). Co-localization of substance P and Met-enkephalin-Arg6-Gly7-Leu8 in the intraspinal neurons of the rat, with special reference to the neurons in the substantia gelatinosa. *Brain Research*, 453(1-2): 110-116.
- Sergeyev, V., Hökfelt, T., and Hurd, Y. (1999). Serotonin and substance P co-exist in dorsal raphe neurons of the human brain. *Clinical Neuroscience and Neuropathology*, 10(18): 3967-3970.
- Seybold, V., and Elde, R. (1980). Immunohistochemical studies of peptidergic neurons in the dorsal horn of the spinal cord. *The Journal of Histochemistry and Cytochemistry*. 28(4): 367-370.
- Simantov, R., Kuhar, M. J., Uhl, G. R., and Snyder, S. H. (1977). Opioid peptide enkephalin: Immunohistochemical mapping in rat central nervous system. *Proceedings of the National Academy of Sciences USA*. 74(5):2167-2171.
- Singer, P. (1990). Do animals feel pain?. *Animal Liberation* (2nd ed.)(pp. 10-15). New York: Avon Books.
- Sommer, C. (2006). Is serotonin hyperalgesic or analgesic?. *Current Pain and Headache Reports*, 10, 101-106.
- Stamford, J. A. (1995). Descending control of pain. *British Journal of Anaesthesia*, 75, 217-227.
- Stensaas, L. J., and Stensaas, S. S. (1967). Astrocytic neuroglial cells, oligodendrocytes and microgliocytes in the spinal cord of the toad. *Cell and Tissue Research*, 84(4): 473-489.
- Stevens, C. W. (2004). Opioid research in amphibians: an alternative pain model yielding insights on the evolution of opioid receptors. *Brain Research Reviews*, 46, 204-215.
- Stuesse S. L., Adli, D. S. H., and Cruce, W. L. R. (2001). Immunohistochemical distribution of enkephalin, substance P and somatostatin in the brainstem of the leopard frog, *Rana pipiens*. *Microscopy Research and Technique*, 54, 229-245.
- Tang, J. M. Y. (2008). *Histological development of selected neural structures of Dark-sided Chorus Frog, Microhyla heymonsi (Amphibia: Anura)*. Thesis of the Bachelor of Science Degree. Division of Biohealth Sciences, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur.

Tang, J. M. Y., Durriyyah S. H. A., and Belabut, D. (2010). Histological development of selected neural structures of dark-sided chorus frog, *Microhyla heymonsi* (Amphibia: Anura). *Malaysian Journal of Science* 29(1): 30-36.

Tashiro, T., and Ruda, M. A. (1988). Immunocytochemical identification of axons containing coexistent serotonin and substance P in the cat lumbar spinal cord. *Peptides*, 9: 383-391.

Tashiro, T., Satoda, T., Matsushima, R., and Mizuno, N. (1990). Distribution of axons showing both enkephalin- and serotonin-like immunoreactivities in the lumbar cord segments of the Japanese monkey (*Macaca fuscata*). *Brain Research*, 512(1):143-6.

Tashiro, T., Takahashi, O., Satoda, T., Matsushima, R., and Mizuno, N. (1987). Immunohistochemical demonstration of coexistence of enkephalin-and substance P-like immunoreactivities in axonal components in the lumbar segments of cat spinal cord. *Brain Research*, 424(2): 391-395.

Tessler, A., Himes, B. T., Krieger, N. R., Murray, M., and Goldberger, M. E. (1985). Sciatic nerve transection produces death of dorsal root ganglion cells and reversible loss of substance P in spinal cord. *Brain Research*, 332(2): 209-218.

Triepel, J., Weindl, A., Kiemle, I., Mader, J., Volz, H. P., Reinecke M., and Forssmann, W. G. (1985). Substance P-immunoreactive neurons in the brainstem of the cat related to cardiovascular centers. *Cell and Tissue Research*, 241(1): 31-41.

Ueda, S., Noiyo, Y., and Sano, Y. (1984). Immunohistochemical demonstration of the serotonin neuron system in the central nervous system of the bullfrog, *Rana catesbeiana*. *Anatomy and Embryology* 169(3): 219-229.

Vecino, E., Piñuela, C., Arévalo, R., Lara, J., Alonso, J. R., and Aijón, J. (1992). Distribution of enkephalin-like immunoreactivity in the central nervous system of the rainbow trout: an immunocytochemical study. *Journal of Anatomy*, 180: 435-453.

Villar, M. J., Wiesenfeld-Hallin, Z., Xu, X. J., Theodorsson, E., Emson, P. C., and Hökfelt, T. (1991). Further studies on galanin-, substance P-, and CGRP-like immunoreactivities in primary sensory neurons and spinal cord: Effects of dorsal rhizotomies and sciatic nerve lesions. *Experimental Neurology*, 112(1): 29-39.

Ville, A. C., Warren, W. F. Jr., and Barnes, R. D. (1978). *General zoology* (5th ed.). Philadelphia: Saunders Company.

Waymire, J. C. (1997). *Biogenic amine neurotransmitters*. Retrieved February 10, 2009 from The University of Texas Health Science Center website: <http://nba.uth.tmc.edu/neuroscience/s1/i12-5.html>

Wessendorf, M. W., Appel, N. M., Molitor, T. W., and Elde, R. P. (1990). A method for immunofluorescent demonstration of three coexisting neurotransmitters in rat brain and spinal cord, using the fluorophores fluorescein, lissamine rhodamine, and 7-amino-4-methylcoumarin-3-acetic acid. *The Journal of Histochemistry and Cytochemistry*, 38(12):1859-77.

Wessendorf, M. W., and Elde, R. (1985). Characterization of an immunofluorescence technique for the demonstration of coexisting neurotransmitters within nerve fibers and terminals. *The Journal of Histochemistry and Cytochemistry*, 33(10): 984-994.

Wessendorf, M. W., and Elde, R. (1987). The coexistence of serotonin- and substance P-like immunoreactivity in the spinal cord of the rat as shown by immunofluorescent double labeling. *The Journal of Neuroscience*, 7(8): 2352-2363.

Wester, K., Wahlund, E., Sundström, C., Ranefall, P., Bengtsson, E., Russell, P. J., et al. (2000). Paraffin section storage and immunohistochemistry: effects of time, temperature, fixation, and retrieval protocol with emphasis on p53 protein and MIB1 antigen. *Applied Immunohistochemistry & Molecular Morphology*, 8(1): 61-70.

Wolters, J. G., Ten Donkelaar, H. J., Steinbusch, H. W. M., and Verhofstad, A. A. J. (1985) Distribution of serotonin in the brain stem and spinal cord of the lizard *Varanus exanthematicus*: An immunohistochemical study. *Neuroscience*, 14(1): 169-193.

Wolters, J. G., Ten Donkelaar, H. J., and Verhofstad, A. A. (1986). Distribution of some peptides (substance P, [Leu]enkephalin, [Met]enkephalin) in the brain stem and spinal cord of a lizard, *Varanus exanthematicus*. *Neuroscience*, 18(4): 917-946.

Xie, R., Chung, J. Y., Ylaya, K., Williams, R. L., Guerrero, N., Nakatsuka, N., et al. (2011). Factors influencing the degradation of archival formalin-fixed paraffin-embedded tissue sections. *Journal of Histochemistry and Cytochemistry*, 59(4): 356-365.

Yamashita, S. (2007). Heat-induced antigen retrieval: Mechanisms and application to histochemistry. *Progress in Histochemistry and Cytochemistry*, 41(3): 141-200.

Zeitz, K. P., Guy, N., Malmberg, A. B., Dirajlal, S., Martin, W. J., Sun, L., et al. (2002). The 5-HT₃ subtype of serotonin receptor contributes to nociceptive processing via a novel subset of myelinated and unmyelinated nociceptors. *The Journal of Neuroscience*, 22(3):1010–1019.

Zhang, E.T., and Craig, A. D. (1997). Morphology and distribution of spinothalamic lamina I neurons in the monkey. *The Journal of Neuroscience*, 17(9): 3274-3284.