

- Abramowicz, M. J., Targovnik, H. M., Varela, V., Cochaux, P., Krawiec, L., Pisarev, M. A., Propato, F. V., Juvenal, G., Chester, H. A., & Vassart, G. (1992). Identification of a mutation in the coding sequence of the human thyroid peroxidase gene causing congenital goiter. *J Clin Invest*, 90(4), 1200-1204.
- Adzhubei, I. A., Schmidt, S., Peshkin, L., Ramensky, V. E., Gerasimova, A., Bork, P., Kondrashov, A. S., & Sunyaev, S. R. (2010). A method and server for predicting damaging missense mutations. *Nat Methods*, 7(4), 248-249.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol*, 215(3), 403-410.
- Alzahrani, A. S., Baitei, E. Y., Zou, M., & Shi, Y. (2006). Clinical case seminar: metastatic follicular thyroid carcinoma arising from congenital goiter as a result of a novel splice donor site mutation in the thyroglobulin gene. *J Clin Endocrinol Metab*, 91(3), 740-746.
- Amar-Singh. (2010). TSH Screening Of Cord Blood In Malaysia – Its Impact (The National Congenital Hypothyroid Screening Programme In Malaysian). [Abstracts]. *The Journal of Endocrinology and Metabolism*, 1(1).
- Avbelj, M., Tahirovic, H., Debeljak, M., Kusekova, M., Toromanovic, A., Krzisnik, C., & Battelino, T. (2007). High prevalence of thyroid peroxidase gene mutations in patients with thyroid dyshormonogenesis. *Eur J Endocrinol*, 156(5), 511-519.
- Bakker, B., Bikker, H., Vulsmma, T., de Randamie, J. S., Wiedijk, B. M., & De Vijlder, J. J. (2000). Two decades of screening for congenital hypothyroidism in The Netherlands: TPO gene mutations in total iodide organification defects (an update). *J Clin Endocrinol Metab*, 85(10), 3708-3712.
- Banga, J. P., Mahadevan, D., Barton, G. J., Sutton, B. J., Saldanha, J. W., Odell, E., & McGregor, A. M. (1990). Prediction of domain organisation and secondary structure of thyroid peroxidase, a human autoantigen involved in destructive thyroiditis. *FEBS Lett*, 266(1-2), 133-141.
- Belfiore, A., La Rosa, G. L., La Porta, G. A., Giuffrida, D., Milazzo, G., Lupo, L., Regalbuto, C., & Vigneri, R. (1992). Cancer risk in patients with cold thyroid nodules: relevance of iodine intake, sex, age, and multinodularity. *Am J Med*, 93(4), 363-369.
- Bernard, G., Poulin, C., Puymirat, J., Sternberg, D., & Shevell, M. (2008). Dosage effect of a dominant CLCN1 mutation: a novel syndrome. *J Child Neurol*, 23(2), 163-166.

Betts, M. J., & Russell, R. B. (2003). Amino acid properties and consequences of substitutions. In M. R. Barnes & I. C. Gray (Eds.), *Bioinformatics for geneticists* (pp. 289-316). Chichester, West Sussex, England ; Hoboken, N.J.: Wiley.

Bikker, H., Baas, F., & De Vijlder, J. J. (1997). Molecular analysis of mutated thyroid peroxidase detected in patients with total iodide organification defects. *J Clin Endocrinol Metab*, 82(2), 649-653.

Bikker, H., Vulsma, T., Baas, F., & de Vijlder, J. J. (1995). Identification of five novel inactivating mutations in the human thyroid peroxidase gene by denaturing gradient gel electrophoresis. *Hum Mutat*, 6(1), 9-16.

Bisi, H., Fernandes, V. S., de Camargo, R. Y., Koch, L., Abdo, A. H., & de Brito, T. (1989). The prevalence of unsuspected thyroid pathology in 300 sequential autopsies, with special reference to the incidental carcinoma. *Cancer*, 64(9), 1888-1893.

Bonert, V., & Friedman, T. C. (2007). The thyroid gland. In T. E. Andreoli, C. C. J. Carpenter & R. L. Cecil (Eds.), *Andreoli and Carpenter's Cecil essentials of medicine* (7th ed., pp. 593-602). Philadelphia: Saunders.

Borroni, B., Grassi, M., Agosti, C., Archetti, S., Costanzi, C., Cornali, C., Caltagirone, C., Caimi, L., Di Luca, M., & Padovani, A. (2006). Cumulative effect of COMT and 5-HTTLPR polymorphisms and their interaction with disease severity and comorbidities on the risk of psychosis in Alzheimer disease. *Am J Geriatr Psychiatry*, 14(4), 343-351.

Brabant, G., Maenhaut, C., Kohrle, J., Scheumann, G., Dralle, H., Hoang-Vu, C., Hesch, R. D., von zur Muhlen, A., Vassart, G., & Dumont, J. E. (1991). Human thyrotropin receptor gene: expression in thyroid tumors and correlation to markers of thyroid differentiation and dedifferentiation. *Mol Cell Endocrinol*, 82(1), R7-12.

Bresson, D., Rebuffat, S. A., & Peraldi-Roux, S. (2005). Localization of the immunodominant region on human thyroid peroxidase in autoimmune thyroid diseases: an update. *J Autoimmune Dis*, 2(1), 2.

Brogna, S., & Wen, J. (2009). Nonsense-mediated mRNA decay (NMD) mechanisms. *Nat Struct Mol Biol*, 16(2), 107-113.

Bronnegard, M., Torring, O., Boos, J., Sylven, C., Marcus, C., & Wallin, G. (1994). Expression of thyrotropin receptor and thyroid hormone receptor messenger ribonucleic acid in normal, hyperplastic, and neoplastic human thyroid tissue. *J Clin Endocrinol Metab*, 79(2), 384-389.

- Buchan, D. W., Ward, S. M., Lobley, A. E., Nugent, T. C., Bryson, K., & Jones, D. T. (2010). Protein annotation and modelling servers at University College London. *Nucleic Acids Res*, 38(Web Server issue), W563-568.
- Caillou, B., Troalen, F., Baudin, E., Talbot, M., Filetti, S., Schlumberger, M., & Bidart, J. M. (1998). Na⁺/I⁻ symporter distribution in human thyroid tissues: an immunohistochemical study. *J Clin Endocrinol Metab*, 83(11), 4102-4106.
- Camargo, R., Limbert, E., Gillam, M., Henriques, M. M., Fernandes, C., Catarino, A. L., Soares, J., Alves, V. A., Kopp, P., & Medeiros-Neto, G. (2001). Aggressive metastatic follicular thyroid carcinoma with anaplastic transformation arising from a long-standing goiter in a patient with Pendred's syndrome. *Thyroid*, 11(10), 981-988.
- Cartegni, L., Chew, S. L., & Krainer, A. R. (2002). Listening to silence and understanding nonsense: exonic mutations that affect splicing. *Nat Rev Genet*, 3(4), 285-298.
- Castro, M. R., Bergert, E. R., Beito, T. G., McIver, B., Goellner, J. R., & Morris, J. C. (1999). Development of monoclonal antibodies against the human sodium iodide symporter: immunohistochemical characterization of this protein in thyroid cells. *J Clin Endocrinol Metab*, 84(8), 2957-2962.
- Cetani, F., Costagliola, S., Tonacchera, M., Panneels, V., Vassart, G., & Ludgate, M. (1995). The thyroperoxidase doublet is not produced by alternative splicing. *Mol Cell Endocrinol*, 115(2), 125-132.
- Chang, Y. F., Imam, J. S., & Wilkinson, M. F. (2007). The nonsense-mediated decay RNA surveillance pathway. *Annu Rev Biochem*, 76, 51-74.
- Charron, M., DeCerbo, J. N., & Wright, W. W. (2003). A GC-box within the proximal promoter region of the rat cathepsin L gene activates transcription in Sertoli cells of sexually mature rats. *Biol Reprod*, 68(5), 1649-1656.
- Chedrese, P. J. (2009). *Reproductive endocrinology : a molecular approach* (1st ed.). New York: Springer.
- Cheron, R. G., Kaplan, M. M., Larsen, P. R., Selenkow, H. A., & Crigler, J. F., Jr. (1981). Neonatal thyroid function after propylthiouracil therapy for maternal Graves' disease. *N Engl J Med*, 304(9), 525-528.
- Chertok Shacham, E., Ishay, A., Irit, E., Pohlenz, J., & Tenenbaum-Rakover, Y. (2012). Minimally invasive follicular thyroid carcinoma developed in dyshormonogenetic multinodular goiter due to thyroid peroxidase gene mutation. *Thyroid*, 22(5), 542-546.

Cheung, V. G., Spielman, R. S., Ewens, K. G., Weber, T. M., Morley, M., & Burdick, J.

T. (2005). Mapping determinants of human gene expression by regional and genome-wide association. *Nature*, 437(7063), 1365-1369.

Coleman, W. B., & Tsongalis, G. J. (2009). *Molecular pathology : the molecular basis of human disease*. Burlington, Mass.: Academic Press.

Cooper, D. S., Axelrod, L., DeGroot, L. J., Vickery, A. L., Jr., & Maloof, F. (1981). Congenital goiter and the development of metastatic follicular carcinoma with evidence for a leak of nonhormonal iodide: clinical, pathological, kinetic, and biochemical studies and a review of the literature. *J Clin Endocrinol Metab*, 52(2), 294-306.

Cooper, D. S., Doherty, G. M., Haugen, B. R., Kloos, R. T., Lee, S. L., Mandel, S. J., Mazzaferri, E. L., McIver, B., Pacini, F., Schlumberger, M., Sherman, S. I., Steward, D. L., & Tuttle, R. M. (2009). Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid*, 19(11), 1167-1214.

D'Avanzo, A., Treseler, P., Ituarte, P. H., Wong, M., Streja, L., Greenspan, F. S., Siperstein, A. E., Duh, Q. Y., & Clark, O. H. (2004). Follicular thyroid carcinoma: histology and prognosis. *Cancer*, 100(6), 1123-1129.

Damante, G., & Di Lauro, R. (1994). Thyroid-specific gene expression. *Biochim Biophys Acta*, 1218(3), 255-266.

de Vijlder, J. J. (2003). Primary congenital hypothyroidism: defects in iodine pathways. *Eur J Endocrinol*, 149(4), 247-256.

de Vijlder, J. J., Ris-Stalpers, C., & Vulsmma, T. (1997). Inborn errors of thyroid hormone biosynthesis. *Exp Clin Endocrinol Diabetes*, 105 Suppl 4, 32-37.

Deladoey, J., Pfarr, N., Vuissoz, J. M., Parma, J., Vassart, G., Biesterfeld, S., Pohlenz, J., & Van Vliet, G. (2008). Pseudodominant inheritance of goitrous congenital hypothyroidism caused by TPO mutations: molecular and in silico studies. *J Clin Endocrinol Metab*, 93(2), 627-633.

Deladoey, J., Vuissoz, J. M., Domene, H. M., Malik, N., Gruneiro-Papendieck, L., Chiesa, A., Heinrich, J. J., & Mullis, P. E. (2003). Congenital secondary hypothyroidism due to a mutation C105Vfs114X thyrotropin-beta mutation: genetic study of five unrelated families from Switzerland and Argentina. *Thyroid*, 13(6), 553-559.

- Delange, F. (1979). Neonatal screening for congenital hypothyroidism in Europe. Report of the Newborn Committee of the European Thyroid Association. *Acta Endocrinol Suppl (Copenh)*, 223, 3-29.
- Delange, F. (1998). Screening for congenital hypothyroidism used as an indicator of the degree of iodine deficiency and of its control. *Thyroid*, 8(12), 1185-1192.
- Delange, F., Dodion, J., Wolter, R., Bourdoux, P., Dalhem, A., Glinoer, D., & Ermans, A. M. (1978). Transient hypothyroidism in the newborn infant. *J Pediatr*, 92(6), 974-976.
- Di Cristofaro, J., Silvy, M., Lanteaume, A., Marcy, M., Carayon, P., & De Micco, C. (2006). Expression of tpo mRNA in thyroid tumors: quantitative PCR analysis and correlation with alterations of ret, Braf , ras and pax8 genes. *Endocr Relat Cancer*, 13(2), 485-495.
- Dunford, H. B. (2001). How do enzymes work? Effect of electron circuits on transition state acid dissociation constants. *J Biol Inorg Chem*, 6(8), 819-822.
- Elisei, R., Vassart, G., & Ludgate, M. (1991). Demonstration of the existence of the alternatively spliced form of thyroid peroxidase in normal thyroid. *J Clin Endocrinol Metab*, 72(3), 700-702.
- Endo, Y., Onogi, S., Umeki, K., Yamamoto, I., Kotani, T., Ohtaki, S., & Fujita, T. (1995). Regional localization of the gene for thyroid peroxidase to human chromosome 2p25 and mouse chromosome 12C. *Genomics*, 25(3), 760-761.
- Eugene, D., Djemli, A., & Van Vliet, G. (2005). Sexual dimorphism of thyroid function in newborns with congenital hypothyroidism. *J Clin Endocrinol Metab*, 90(5), 2696-2700.
- Ferrand, M., Le Fourn, V., & Franc, J. L. (2003). Increasing diversity of human thyroperoxidase generated by alternative splicing. Characterized by molecular cloning of new transcripts with single- and multisPLICED mRNAs. *J Biol Chem*, 278(6), 3793-3800.
- Finke, R., Seto, P., Ruf, J., Carayon, P., & Rapoport, B. (1991). Determination at the molecular level of a B-cell epitope on thyroid peroxidase likely to be associated with autoimmune thyroid disease. *J Clin Endocrinol Metab*, 73(4), 919-921.
- Finley, D. J., Zhu, B., Barden, C. B., & Fahey, T. J., 3rd. (2004). Discrimination of benign and malignant thyroid nodules by molecular profiling. *Ann Surg*, 240(3), 425-436; discussion 436-427.

- Fisher, D. A., Dussault, J. H., Foley, T. P., Jr., Klein, A. H., LaFranchi, S., Larsen, P. R., Mitchell, M. L., Murphey, W. H., & Walfish, P. G. (1979). Screening for congenital hypothyroidism: results of screening one million North American infants. *J Pediatr*, 94(5), 700-705.
- Frischmeyer, P. A., & Dietz, H. C. (1999). Nonsense-mediated mRNA decay in health and disease. *Hum Mol Genet*, 8(10), 1893-1900.
- Fruzza, A. G., Demeterco-Berggren, C., & Jones, K. L. (2012). Unawareness of the effects of soy intake on the management of congenital hypothyroidism. *Pediatrics*, 130(3), e699-702.
- Fugazzola, L., Cerutti, N., Mannavola, D., Vannucchi, G., Fallini, C., Persani, L., & Beck-Peccoz, P. (2003). Monoallelic expression of mutant thyroid peroxidase allele causing total iodide organification defect. *J Clin Endocrinol Metab*, 88(7), 3264-3271.
- Gardas, A., Lewartowska, A., Sutton, B. J., Pasieka, Z., McGregor, A. M., & Banga, J. P. (1997). Human thyroid peroxidase (TPO) isoforms, TPO-1 and TPO-2: analysis of protein expression in Graves' thyroid tissue. *J Clin Endocrinol Metab*, 82(11), 3752-3757.
- Granner, D. K. (1999). Thyroid hormones *Harper's biochemistry* (25 ed., pp. 561-566). US: McGraw-Hill Publishing Co.
- Gruters, A. (1992). Congenital hypothyroidism. *Pediatr Ann*, 21(1), 15, 18-21, 24-18.
- Gruters, A., & Krude, H. (2007). Update on the management of congenital hypothyroidism. *Horm Res*, 68 Suppl 5, 107-111.
- Gruters, A., Krude, H., & Biebermann, H. (2004). Molecular genetic defects in congenital hypothyroidism. *Eur J Endocrinol*, 151 Suppl 3, U39-44.
- Han, W., Kim, K. Y., Yang, S. J., Noh, D. Y., Kang, D., & Kwack, K. (2012). SNP-SNP interactions between DNA repair genes were associated with breast cancer risk in a Korean population. *Cancer*, 118(3), 594-602.
- Hapgood, J. P., Riedemann, J., & Scherer, S. D. (2001). Regulation of gene expression by GC-rich DNA cis-elements. *Cell Biol Int*, 25(1), 17-31.
- Hawthorn, L., Stein, L., Varma, R., Wiseman, S., Loree, T., & Tan, D. (2004). TIMP1 and SERPIN-A overexpression and TFF3 and CRABP1 underexpression as biomarkers for papillary thyroid carcinoma. *Head Neck*, 26(12), 1069-1083.

- Hendry, E., Taylor, G., Ziemnicka, K., Grennan Jones, F., Furmaniak, J., & Rees Smith, B. (1999). Recombinant human thyroid peroxidase expressed in insect cells is soluble at high concentrations and forms diffracting crystals. *J Endocrinol*, 160(3), R13-15.
- Hobby, P., Gardas, A., Radomski, R., McGregor, A. M., Banga, J. P., & Sutton, B. J. (2000). Identification of an immunodominant region recognized by human autoantibodies in a three-dimensional model of thyroid peroxidase. *Endocrinology*, 141(6), 2018-2026.
- Hoyes, A. D., & Kershaw, D. R. (1985). Anatomy and development of the thyroid gland. *Ear Nose Throat J*, 64(7), 318-333.
- Hsiau, A. c. (2000). *Contemporary Taiwanese cultural nationalism*. London ; New York: Routledge.
- Huang, Y., Prasad, M., Lemon, W. J., Hampel, H., Wright, F. A., Kornacker, K., LiVolsi, V., Frankel, W., Kloos, R. T., Eng, C., Pellegata, N. S., & de la Chapelle, A. (2001). Gene expression in papillary thyroid carcinoma reveals highly consistent profiles. *Proc Natl Acad Sci U S A*, 98(26), 15044-15049.
- Hulse, J. A. (1984). Outcome for congenital hypothyroidism. *Arch Dis Child*, 59(1), 23-29.
- Ieiri, T., Cochaux, P., Targovnik, H. M., Suzuki, M., Shimoda, S., Perret, J., & Vassart, G. (1991). A 3' splice site mutation in the thyroglobulin gene responsible for congenital goiter with hypothyroidism. *J Clin Invest*, 88(6), 1901-1905.
- Jhiang, S. M., Cho, J. Y., Ryu, K. Y., DeYoung, B. R., Smanik, P. A., McGaughy, V. R., Fischer, A. H., & Mazzaferri, E. L. (1998). An immunohistochemical study of Na⁺/I⁻ symporter in human thyroid tissues and salivary gland tissues. *Endocrinology*, 139(10), 4416-4419.
- Joba, W., Spitzweg, C., Schriever, K., & Heufelder, A. E. (1999). Analysis of human sodium/iodide symporter, thyroid transcription factor-1, and paired-box-protein-8 gene expression in benign thyroid diseases. *Thyroid*, 9(5), 455-466.
- Kalina, M., Weavers, B., & Pearse, A. G. (1971). Ultrastructural localization of succinate dehydrogenase in mouse liver mitochondria; a cytochemical study. *J Histochem Cytochem*, 19(2), 124-130.
- Kang, I. N. (2010). *Functional characterisation of a novel N132D mutation in the thyroid transcription factor-2 (TTF-2) gene and mutational analysis of the thyroid peroxidase (TPO) gene in patients with congenital hypothyroidism*. Master in Medical Science, University Of Malaya, Kuala Lumpur. (150)

Karger, S., Engelhardt, C., Eszlinger, M., Tonjes, A., Herrmann, F., Muller, P., Schmidt, T., Weiss, C. L., Dralle, H., Lippitzsch, F., Tannapfel, A., & Fuhrer, D. (2006). Cytology and mRNA expression analysis of fine needle aspirates of thyroid nodules in an East German region with borderline iodine deficiency. *Horm Metab Res*, 38(10), 662-667.

Kelly, M., & Semsarian, C. (2009). Multiple mutations in genetic cardiovascular disease: a marker of disease severity? *Circ Cardiovasc Genet*, 2(2), 182-190.

Kimura, S., Hong, Y. S., Kotani, T., Ohtaki, S., & Kikkawa, F. (1989). Structure of the human thyroid peroxidase gene: comparison and relationship to the human myeloperoxidase gene. *Biochemistry*, 28(10), 4481-4489.

Kimura, S., Kotani, T., McBride, O. W., Umeki, K., Hirai, K., Nakayama, T., & Ohtaki, S. (1987). Human thyroid peroxidase: complete cDNA and protein sequence, chromosome mapping, and identification of two alternately spliced mRNAs. *Proc Natl Acad Sci U S A*, 84(16), 5555-5559.

Lagorce, J. F., Thomes, J. C., Catanzano, G., Buxeraud, J., Raby, M., & Raby, C. (1991). Formation of molecular iodine during oxidation of iodide by the peroxidase/H₂O₂ system. Implications for antithyroid therapy. *Biochem Pharmacol*, 42 Suppl, S89-92.

Lai, O. F., Zaiden, N., Goh, S. S., Mohamed, N. E., Seah, L. L., Fong, K. S., Estienne, V., Carayon, P., Ho, S. C., & Khoo, D. H. (2006). Detection of thyroid peroxidase mRNA and protein in orbital tissue. *Eur J Endocrinol*, 155(2), 213-218.

Lalonde, E., Ha, K. C., Wang, Z., Bemmo, A., Kleinman, C. L., Kwan, T., Pastinen, T., & Majewski, J. (2011). RNA sequencing reveals the role of splicing polymorphisms in regulating human gene expression. *Genome Res*, 21(4), 545-554.

Lazar, V., Bidart, J. M., Caillou, B., Mahe, C., Lacroix, L., Filetti, S., & Schlumberger, M. (1999). Expression of the Na⁺/I⁻ symporter gene in human thyroid tumors: a comparison study with other thyroid-specific genes. *J Clin Endocrinol Metab*, 84(9), 3228-3234.

Lee, K. H., & Tan, C. B. (1999). *The Chinese in Malaysia*. New York: Oxford University Press.

Lee, S. T., Lee, D. H., Kim, J. Y., Kwon, M. J., Kim, J. W., Hong, Y. H., Lee, Y. W., & Ki, C. S. (2011). Molecular screening of the TSH receptor (TSHR) and thyroid peroxidase (TPO) genes in Korean patients with nonsyndromic congenital hypothyroidism. *Clin Endocrinol (Oxf)*, 75(5), 715-721.

Levy, O., Dai, G., Riedel, C., Ginter, C. S., Paul, E. M., Lebowitz, A. N., & Carrasco, N. (1997). Characterization of the thyroid Na⁺/I⁻ symporter with an anti-COOH terminus antibody. *Proc Natl Acad Sci U S A*, 94(11), 5568-5573.

Li, Q., Fallin, M. D., Louis, T. A., Lasseter, V. K., McGrath, J. A., Avramopoulos, D., Wolyniec, P. S., Valle, D., Liang, K. Y., Pulver, A. E., & Ruczinski, I. (2010). Detection of SNP-SNP interactions in trios of parents with schizophrenic children. *Genet Epidemiol*, 34(5), 396-406.

Libert, F., Ruel, J., Ludgate, M., Swillens, S., Alexander, N., Vassart, G., & Dinsart, C. (1987). Thyroperoxidase, an auto-antigen with a mosaic structure made of nuclear and mitochondrial gene modules. *EMBO J*, 6(13), 4193-4196.

Lin, C. P., Chen, W., & Wu, K. W. (1994). Povidone-iodine in umbilical cord care interferes with neonatal screening for hypothyroidism. *Eur J Pediatr*, 153(10), 756-758.

Liu, C. B., Wang, F. J., Wu, C. J., & Zhou, M. (2010). TPO and DIO1 Mutations in Patients With the Coexistence of Hashimoto Thyroiditis and Papillary Thyroid Carcinoma. *Endocrinologist*, 20(5), 254-258.

Lv, B., Tang, Y., Li, X., Wang, G., Yuan, C., Liu, Y., Jiang, L., & Xiao, X. (2009). Identification and characterization of the minimal promoter of Mipu1: the role of GC boxes in the regulation of basal transcription. *Acta Biochim Biophys Sin (Shanghai)*, 41(4), 309-315.

Macchia, P. E. (2000). Recent advances in understanding the molecular basis of primary congenital hypothyroidism. *Mol Med Today*, 6(1), 36-42.

Mackenzie, E. J., & Mortimer, R. H. (2004). 6: Thyroid nodules and thyroid cancer. *Med J Aust*, 180(5), 242-247.

Magnusson, R. P., Chazenbalk, G. D., Gestautas, J., Seto, P., Filetti, S., DeGroot, L. J., & Rapoport, B. (1987). Molecular cloning of the complementary deoxyribonucleic acid for human thyroid peroxidase. *Mol Endocrinol*, 1(11), 856-861.

Magnusson, R. P., Taurog, A., & Dorris, M. L. (1984a). Mechanism of iodide-dependent catalytic activity of thyroid peroxidase and lactoperoxidase. *J Biol Chem*, 259(1), 197-205.

Magnusson, R. P., Taurog, A., & Dorris, M. L. (1984b). Mechanisms of thyroid peroxidase- and lactoperoxidase-catalyzed reactions involving iodide. *J Biol Chem*, 259(22), 13783-13790.

- Mangklabruks, A., Billerbeck, A. E., Wajchenberg, B., Knobel, M., Cox, N. J., DeGroot, L. J., & Medeiros-Neto, G. (1991). Genetic linkage studies of thyroid peroxidase (TPO) gene in families with TPO deficiency. *J Clin Endocrinol Metab*, 72(2), 471-476.
- Maquat, L. E. (2005). Nonsense-mediated mRNA decay in mammals. *J Cell Sci*, 118(Pt 9), 1773-1776.
- Marino, M., & McCluskey, R. T. (2000). Role of thyroglobulin endocytic pathways in the control of thyroid hormone release. *Am J Physiol Cell Physiol*, 279(5), C1295-1306.
- McDonald, D. O., & Pearce, S. H. (2009). Thyroid peroxidase forms thionamide-sensitive homodimers: relevance for immunomodulation of thyroid autoimmunity. *J Mol Med (Berl)*, 87(10), 971-980.
- McLachlan, S. M., & Rapoport, B. (1992). The molecular biology of thyroid peroxidase: cloning, expression and role as autoantigen in autoimmune thyroid disease. *Endocr Rev*, 13(2), 192-206.
- Medeiros-Neto, G., Gil-Da-Costa, M. J., Santos, C. L., Medina, A. M., Silva, J. C., Tsou, R. M., & Sobrinho-Simoes, M. (1998). Metastatic thyroid carcinoma arising from congenital goiter due to mutation in the thyroperoxidase gene. *J Clin Endocrinol Metab*, 83(11), 4162-4166.
- Medeiros-Neto, G., Targovnik, H. M., & Vassart, G. (1993). Defective thyroglobulin synthesis and secretion causing goiter and hypothyroidism. *Endocr Rev*, 14(2), 165-183.
- Mount, D. W. (2007). Using the Basic Local Alignment Search Tool (BLAST). *CSH Protoc*, 2007, pdb top17.
- Musa, M., Harun, F., & Junit, S. M. (2006a). Mutational analysis of genes encoding thyroid stimulating hormone receptor (TSHR) and thyroid transcription factor, FOXE1 in patients with congenital hypothyroidism. *Mal J Biochem Mol Biol*, 15(1), 8-18.
- Musa, M., Harun, F., & Junit, S. M. (2006b). Mutational analysis of genes encoding thyroid stimulating hormone receptor (TSHR) and thyroid transcription factor, FOXE1 in patients with congenital hypothyroidism. *Malaysian Journal of Biochemistry and Molecular Biology*, 15.
- Ng, P. C., & Henikoff, S. (2001). Predicting deleterious amino acid substitutions. *Genome Res*, 11(5), 863-874.

- Niu, D. M., Hwang, B., Chu, Y. K., Liao, C. J., Wang, P. L., & Lin, C. Y. (2002). High prevalence of a novel mutation (2268 insT) of the thyroid peroxidase gene in Taiwanese patients with total iodide organification defect, and evidence for a founder effect. *J Clin Endocrinol Metab*, 87(9), 4208-4212.
- Niu, D. M., Lin, C. Y., Hwang, B., Jap, T. S., Liao, C. J., & Wu, J. Y. (2005). Contribution of genetic factors to neonatal transient hypothyroidism. *Arch Dis Child Fetal Neonatal Ed*, 90(1), F69-72.
- Nordlund, T. M. (2011). *Quantitative understanding of biosystems : an introduction to biophysics*. Boca Raton: Taylor & Francis.
- Nugent, T., & Jones, D. T. (2009). Transmembrane protein topology prediction using support vector machines. *BMC Bioinformatics*, 10, 159.
- O'Shea, P. J., Harvey, C. B., Suzuki, H., Kaneshige, M., Kaneshige, K., Cheng, S. Y., & Williams, G. R. (2003). A thyrotoxic skeletal phenotype of advanced bone formation in mice with resistance to thyroid hormone. *Mol Endocrinol*, 17(7), 1410-1424.
- Ohta, K., Endo, T., & Onaya, T. (1991). The mRNA levels of thyrotropin receptor, thyroglobulin and thyroid peroxidase in neoplastic human thyroid tissues. *Biochem Biophys Res Commun*, 174(3), 1148-1153.
- Ordoonkhani, A., Mirmiran, P., Walfish, P. G., & Azizi, F. (2007). Transient neonatal hypothyroidism is associated with elevated serum anti-thyroglobulin antibody levels in newborns and their mothers. *J Pediatr*, 150(3), 315-317, 317 e312.
- Pannain, S., Weiss, R. E., Jackson, C. E., Dian, D., Beck, J. C., Sheffield, V. C., Cox, N., & Refetoff, S. (1999). Two different mutations in the thyroid peroxidase gene of a large inbred Amish kindred: power and limits of homozygosity mapping. *J Clin Endocrinol Metab*, 84(3), 1061-1071.
- Park, S. M., & Chatterjee, V. K. (2005). *Genetics of congenital hypothyroidism* (2005/05/03 ed. Vol. 42).
- Passon, N., Puppin, C., Lavarone, E., Bregant, E., Franzoni, A., Hershman, J., Fenton, M., D'Agostino, M., Durante, C., Russo, D., Filetti, S., & Damante, G. (2012). CREM inhibits promoter activity of NIS gene in thyroid cancer cells. *Thyroid*.
- Pauws, E., Veenboer, G. J., Smit, J. W., de Vijlder, J. J., Morreau, H., & Ris-Stalpers, C. (2004). Genes differentially expressed in thyroid carcinoma identified by comparison of SAGE expression profiles. *FASEB J*, 18(3), 560-561.

Perez-Cuvit, E., Crigler, J. F., Jr., & Stanbury, J. B. (1977). Partial and total iodide organification defect in different sibships in a kindred. *Am J Hum Genet*, 29(2), 142-148.

Pezzini, A., Grassi, M., Del Zotto, E., Archetti, S., Spezi, R., Vergani, V., Assanelli, D., Caimi, L., & Padovani, A. (2005). Cumulative effect of predisposing genotypes and their interaction with modifiable factors on the risk of ischemic stroke in young adults. *Stroke*, 36(3), 533-539.

Phillips, M. A., Fletterick, R., & Rutter, W. J. (1990). Arginine 127 stabilizes the transition state in carboxypeptidase. *J Biol Chem*, 265(33), 20692-20698.

Piippo, K., Swan, H., Pasternack, M., Chapman, H., Paavonen, K., Viitasalo, M., Toivonen, L., & Kontula, K. (2001). A founder mutation of the potassium channel KCNQ1 in long QT syndrome: implications for estimation of disease prevalence and molecular diagnostics. *J Am Coll Cardiol*, 37(2), 562-568.

Pohlenz, J., & Refetoff, S. (1999). Mutations in the sodium/iodide symporter (NIS) gene as a cause for iodide transport defects and congenital hypothyroidism. *Biochimie*, 81(5), 469-476.

Prasad, N. B., Somervell, H., Tufano, R. P., Dackiw, A. P., Marohn, M. R., Califano, J. A., Wang, Y., Westra, W. H., Clark, D. P., Umbrecht, C. B., Libutti, S. K., & Zeiger, M. A. (2008). Identification of genes differentially expressed in benign versus malignant thyroid tumors. *Clin Cancer Res*, 14(11), 3327-3337.

Rastogi, M. V., & LaFranchi, S. H. (2010). Congenital hypothyroidism. *Orphanet J Rare Dis*, 5, 17.

Ricketts, M. H., Simons, M. J., Parma, J., Mercken, L., Dong, Q., & Vassart, G. (1987). A nonsense mutation causes hereditary goitre in the Afrikander cattle and unmasks alternative splicing of thyroglobulin transcripts. *Proc Natl Acad Sci U S A*, 84(10), 3181-3184.

Ringel, M. D., Anderson, J., Souza, S. L., Burch, H. B., Tambascia, M., Shriver, C. D., & Tuttle, R. M. (2001). Expression of the sodium iodide symporter and thyroglobulin genes are reduced in papillary thyroid cancer. *Mod Pathol*, 14(4), 289-296.

Ris-Stalpers, C., & Bikker, H. (2010). Genetics and phenomics of hypothyroidism and goiter due to TPO mutations. *Mol Cell Endocrinol*, 322(1-2), 38-43.

- Rivolta, C. M., Esperante, S. A., Gruneiro-Papendieck, L., Chiesa, A., Moya, C. M., Domene, S., Varela, V., & Targovnik, H. M. (2003). Five novel inactivating mutations in the thyroid peroxidase gene responsible for congenital goiter and iodide organification defect. *Hum Mutat*, 22(3), 259.
- Roberts, C. G., & Ladenson, P. W. (2004). Hypothyroidism. *Lancet*, 363(9411), 793-803.
- Rosai, J., Carcangiu, M. L., & DeLellis, R. A. (1991). Tumors of the thyroid gland. In: *Atlas of tumor pathology. Third series* (pp. 21-48). Washington: Armed Forces Institute of Pathology.
- Rose, S. R., Brown, R. S., Foley, T., Kaplowitz, P. B., Kaye, C. I., Sundararajan, S., & Varma, S. K. (2006). Update of newborn screening and therapy for congenital hypothyroidism. *Pediatrics*, 117(6), 2290-2303.
- Ross, D. S. (2001). Serum thyroid-stimulating hormone measurement for assessment of thyroid function and disease. *Endocrinol Metab Clin North Am*, 30(2), 245-264, vii.
- Ruf, J., & Carayon, P. (2006). Structural and functional aspects of thyroid peroxidase. *Arch Biochem Biophys*, 445(2), 269-277.
- Saito, T., Endo, T., Kawaguchi, A., Ikeda, M., Katoh, R., Kawaoi, A., Muramatsu, A., & Onaya, T. (1998). Increased expression of the sodium/iodide symporter in papillary thyroid carcinomas. *J Clin Invest*, 101(7), 1296-1300.
- Sali, A., & Blundell, T. L. (1993). Comparative protein modelling by satisfaction of spatial restraints. *J Mol Biol*, 234(3), 779-815.
- Sapini Y, R. P., Nor Zuraida Z. (2010). Thyroid disorders and psychiatric morbidities. *MJP Online Early*, MJP 02-08-09.
- Sarkar, S. D., & Becker, D. V. (2001). Thyroid uptake and imaging. In K. L. Becker (Ed.), *Principles and practice of endocrinology and metabolism* (3rd ed., pp. 336-342). Philadelphia: Lippincott Williams and Wilkins.
- Sethuraman, G., Sugandhan, S., Sharma, G., Chandramohan, K., Chandra, N. C., Dash, S. S., Komal, A., & Sharma, V. K. (2007). Familial homozygous hypercholesterolemia: report of two patients and review of the literature. *Pediatr Dermatol*, 24(3), 230-234.

- Seto, P., Hirayu, H., Magnusson, R. P., Gestautas, J., Portmann, L., DeGroot, L. J., & Rapoport, B. (1987). Isolation of a complementary DNA clone for thyroid microsomal antigen. Homology with the gene for thyroid peroxidase. *J Clin Invest*, 80(4), 1205-1208.
- Sheils, O. M., & Sweeney, E. C. (1999). TSH receptor status of thyroid neoplasms--TaqMan RT-PCR analysis of archival material. *J Pathol*, 188(1), 87-92.
- Sherwood, L., & Cengage Learning (Firm). (2010). *Human physiology : from cells to systems* (7th ed.). Australia ; United States: Brooks/Cole, Cengage Learning.
- Silverberg, S. G., & Vidone, R. A. (1966). Adenoma and carcinoma of the thyroid. *Cancer*, 19(8), 1053-1062.
- Singh, M., Singh, P., Juneja, P. K., Singh, S., & Kaur, T. (2011). SNP-SNP interactions within APOE gene influence plasma lipids in postmenopausal osteoporosis. *Rheumatol Int*, 31(3), 421-423.
- Sodre, A. K., Rubio, I. G., Galrao, A. L., Knobel, M., Tomimori, E. K., Alves, V. A., Kanamura, C. T., Buchpiguel, C. A., Watanabe, T., Friguglietti, C. U., Kulcsar, M. A., Medeiros-Neto, G., & Camargo, R. Y. (2008). Association of low sodium-iodide symporter messenger ribonucleic acid expression in malignant thyroid nodules with increased intracellular protein staining. *J Clin Endocrinol Metab*, 93(10), 4141-4145.
- Spencer, C. A., Takeuchi, M., & Kazarosyan, M. (1996). Current status and performance goals for serum thyrotropin (TSH) assays. *Clin Chem*, 42(1), 140-145.
- Sterne-Weiler, T., Howard, J., Mort, M., Cooper, D. N., & Sanford, J. R. (2011). Loss of exon identity is a common mechanism of human inherited disease. *Genome Res*, 21(10), 1563-1571.
- Stevens, T. M., Richards, A. T., Bewtra, C., & Sharma, P. (2011). Tumors metastatic to thyroid neoplasms: a case report and review of the literature. *Patholog Res Int*, 2011, 238693.
- Summers, J. E. (1950). Surgical anatomy of the thyroid gland. *Am J Surg*, 80(1), 35-43.
- Sun, X., & Maquat, L. E. (2000). mRNA surveillance in mammalian cells: the relationship between introns and translation termination. *RNA*, 6(1), 1-8.

- Takasu, N., Ohno, S., Komiya, I., & Yamada, T. (1992). Requirements of follicle structure for thyroid hormone synthesis; cytoskeletons and iodine metabolism in polarized monolayer cells on collagen gel and in double layered, follicle-forming cells. *Endocrinology*, 131(3), 1143-1148.
- Taurog, A., & Wall, M. (1998). Proximal and distal histidines in thyroid peroxidase: relation to the alternatively spliced form, TPO-2. *Thyroid*, 8(2), 185-191.
- Toft, A. D. (1994). Thyroxine therapy. *N Engl J Med*, 331(3), 174-180.
- Tsunoda, T., & Takagi, T. (1999). Estimating transcription factor bindability on DNA. *Bioinformatics*, 15(7-8), 622-630.
- Umeki, K., Yamamoto, I., Yukizane, S., & Kotani, T. (2004). Congenital hypothyroidism caused by a unique thyroid peroxidase allele containing two mutations, C1708T and C2737T. *J Pediatr Endocrinol Metab*, 17(2), 231-234.
- Valentine, C. R. (1998). The association of nonsense codons with exon skipping. *Mutat Res*, 411(2), 87-117.
- van de Graaf, S. A., Ris-Stalpers, C., Pauws, E., Mendive, F. M., Targovnik, H. M., & de Vijlder, J. J. (2001). Up to date with human thyroglobulin. *J Endocrinol*, 170(2), 307-321.
- Van Vliet, G. (2003). Development of the thyroid gland: lessons from congenitally hypothyroid mice and men. *Clin Genet*, 63(6), 445-455.
- Varela, V., Rivolta, C. M., Esperante, S. A., Gruneiro-Papendieck, L., Chiesa, A., & Targovnik, H. M. (2006). Three mutations (p.Q36H, p.G418fsX482, and g.IVS19-2A>C) in the dual oxidase 2 gene responsible for congenital goiter and iodide organification defect. *Clin Chem*, 52(2), 182-191.
- Vogelsang, M., Wang, Y., Veber, N., Mwapagha, L. M., & Parker, M. I. (2012). The cumulative effects of polymorphisms in the DNA mismatch repair genes and tobacco smoking in oesophageal cancer risk. *PLoS One*, 7(5), e36962.
- Wallner, B., & Elofsson, A. (2003). Can correct protein models be identified? *Protein Sci*, 12(5), 1073-1086.
- Warf, M. B., & Berglund, J. A. (2010). Role of RNA structure in regulating pre-mRNA splicing. *Trends Biochem Sci*, 35(3), 169-178.

- Weischenfeldt, J., Lykke-Andersen, J., & Porse, B. (2005). Messenger RNA surveillance: neutralizing natural nonsense. *Curr Biol*, 15(14), R559-562.
- Whitley, R. J. (1999). Thyroid function. In C. A. Burtis, E. R. Ashwood & N. W. Tietz (Eds.), *Tietz textbook of clinical chemistry* (3rd ed., pp. 1496-1529). Philadelphia: W.B. Saunders.
- Wu, J. Y., Shu, S. G., Yang, C. F., Lee, C. C., & Tsai, F. J. (2002). Mutation analysis of thyroid peroxidase gene in Chinese patients with total iodide organification defect: identification of five novel mutations. *J Endocrinol*, 172(3), 627-635.
- Yen, P. M. (2001). Physiological and molecular basis of thyroid hormone action. *Physiol Rev*, 81(3), 1097-1142.
- Yong, P. H., Harun, F., & Junit, S. M. (2007). Molecular analysis of PAX8 gene in unrelated patients with congenital hypothyroidism *The Internet J Endocrinol*, 3(2).
- Zanelli, E., Henry, M., Charvet, B., & Malthiery, Y. (1990). Evidence for an alternate splicing in the thyroperoxidase messenger from patients with Graves' disease. *Biochem Biophys Res Commun*, 170(2), 735-741.
- Zhao, J., Hyman, L., & Moore, C. (1999). Formation of mRNA 3' ends in eukaryotes: mechanism, regulation, and interrelationships with other steps in mRNA synthesis. *Microbiol Mol Biol Rev*, 63(2), 405-445.