

REFERENCES

References

- Abdul Jalil Muhamad and Mohd Jamaal Hoesni (1992). Possible source for Tembungo oils: evidences from biomarker fingerprints. *Bulletin of the Geological Society of Malaysia*, 32, 213-232.
- Albaghdady, A., Wan Hasiah, Abdullah and Lee Chai Peng (2003). An organic geochemical study of the Miocene sedimentary sequence of Labuan Island, offshore western Sabah, East Malaysia. Annual Geological Conference 2003, May 24-26, Kuching, Sarawak, Malaysia. *Bulletin of the Geological Society of Malaysia*, 46, 455-460
- Alberdi, M., Lopez, L. (2000) Biomarker 18(H)-oleanane: a geochemical tool to assess Venezuelan petroleum systems. *Journal of South American Earth Sciences*. 13, 751-759.
- Alexander, R., Kagi, R., and Noble, R. (1981). Geochemical correlation of Windalia oil and extracts of Winning Group (Cretaceous) potential source rocks, Barrow sub basin, Western Australia. *American Association of Petroleum Geologists Bulletin*, 65, 235-250.
- Azrina Anuar and Abdul Jalil Muhamad (1997). A comparison of Source rock facies and hydrocarbon types of the Middle Miocene sequence, offshore NW Sabah Basin, Malaysia. In: Howes, J. V. C. and Noble, R. A.,eds., Proceedings of the IPA Petroleum Systems of SE Asia and Australia, 21-23 May 1997. *Indonesian Petroleum Association*, Jakarta, 773-786.
- Azrina Anuar and Kinghorn, R. R. F. (1994) Sterane and triterpane biomarker characteristic from oils and sediment extracts of the Middle-Upper Miocene sequence, North Sabah basin. AAPG International conference and exhibition, August 21-24, 1994. *Bulletin of the Geological Society of Malaysia*, 37, 415-436
- Beddoes, L. R., Jr. (1976) The balabac Sub-Basin, southwestern Sulu Sea, Philippines. Offshore South East Asia Conference, *Seapax Program*, Feb 1976. Paper 15, 22P.
- Blumer, M Mullin, M. M. and Thomas, D. W. (1963) Pristane in zooplankton. *Science*, 140, 974.
- Bol, A. J. and Van Hoorn, B., (1980) Structural styles in western Sabah offshore. *Bulletin of the Geological Society of Malaysia*, 12, 1-16.
- Brassell S. C. and Eglinton G. (1986) Molecular geochemistry indicators in sediments. In Organic Marine geochemistry (Edited by Sohn M.), p. 10- 32. *American Chemical Society*, Washington, D. C.
- Bray, E.E. and Evans, E.D. (1961) Distribution of n-paraffins as a clue to recognition of source beds. *Geochimica et Cosmochimica Acta*, 22, 2-15.
- Brooks, J. D., Gould, K., and Smith, J. W. (1969) Isopernoid hydrocarbons in coal and petroleum. *Nature*, 222, 257-259

- Connan, Bourouille, J., Dessort, D. and Albrecht, P. (1986). The microbial input in carbonate-anhydrite facies of asabkha paleoenvironment from Guatemala: a molecular approach. In: Advance in organic Geochemistry 1985, D. Leythaeuser and J. Rullkötter (Eds.). *Pergamon Press. Oxford*, 29-50
- Connan, J., Cassou, A. M., (1980) Properties of gases and petroleum liquids derived from terrestrial Kerogen at various maturation levels. *Geochimica et Cosmochimica Acta*, 44, 1-23.
- Cook A. C., and Stuckmayer, H. (1986). The role of Coal as a Source Rock for oil. In: Second Southeastern Australia Oil Exploration Symposium. *Glierie, R. C. Ed, Melbourne*, 419-432
- Cooper, B. S. (1990) Practical petroleum geochemistry, *London W4 4PH*, 174 p
- Didyk, B. H., Simoneit, B. R. T., Brassell, S. C., Eglinton, G. (1978) Organic geochemical indicators of paleoenvironmental conditions of sedimentation. *Nature* 272, 216-221
- Eglinton, G. and Calvin, M. (1967) Chemical fossils. *Scientific American*, 216, 32-43.
- Ekweozor, C.M., Udo, O.T., (1987). The oleananes: Origin, maturation and limits of occurrence in southern Nigeria sedimentary basin. Matavelli, L., Novelli, L. (Eds.). *Advances in Organic Geochemistry* 13, 131-141.
- Farrimond, p., Taylor, A., Telnaes, N., (1998) Biomarker maturity parameters: the role of generation and thermal degradation, *organic geochemistry*, 29, 5-7, 1181-1197.
- Fu Jiamo, Sheng Guoying, Xu Jiayou, Eglinton, G., Gowar, A. P., Jia Rongfen, Fan Shanfa, Peng pingan, 1990) Application of biological markers in the assessment of paleoenvironments of Chinese non- marine sediments. *Organic geochemistry*, 16, 4-6, 769-779.
- Graham, U. M. and Robertson, J. D (1995). Micro-PIXE analysis of framboidal pyrite and associated types in oil shale, *Fuel*, 74, No. 4, 530 -535
- Grantham, P. J. (1986) Sterane isomerization and moretane/ hopane ratios in crude oils derived from Tertiary source rocks. *Organic Geochemistry*, Vol. 9, PP. 293-304.
- Hamilton, W. (1979). Tectonics of the Indonesian Region. *U. S. Geol. Surv. Professional Paper* 1078 345 P.
- ten Haven H. L., de Leeuw J. W., Peakman, T. M. and Maxwell, J. R. (1986) Anomalies in steroid and hopanoid maturity indices. *Geochimica et Cosmochimica Acta*, 50, 853-855.
- ten Haven H. L., de Leeuw J. W., Rullkötter J. Sinninghe-Damste J. S. (1987) Restricted utility of pristane / Phytane ratio as Paleoenvironmental indicator. *Nature* 330, pp 641-643

- ten Haven H. L., Rechka, J. R., de Leeuw J. W. and Maxwell, J. R. (1989) Occurrence of (20R)- and (20S)- $\Delta^{8(14)}$ and Δ^{14} 5 α (H)-steranes and the origin of 5 α (H), 14 β (H), 17 β (H)-steranes in an immature sediment. *Geochimica et Cosmochimica Acta*, 53, 2001-2009.
- ten Haven H. L., Rechka, J. R., de Leeuw J. W. and Maxwell, J. R. (1989) Occurrence of (20R) - and (20S) - $\Delta^{8(14)}$ and Δ^{14} 5 α (H)-steranes and the origin of 5 α (H), 14 β (H), 17 β (H)-steranes in an immature sediment. *Geochimica et Cosmochimica Acta*, 53, 2001-2009.
- ten Haven, H. L., Rulkotter, J., (1988). The diagenetic fate of taraxer-14-ene and oleanane isomers. *Geochimica et Cosmochimica Acta*, 56, 2001-2024.
- Hazebroek, H. P. and Tan, D. N. K. (1993) Tertiary tectonic evolution of the NW Sabah continental margin. In proc. Symp. Tectonic Framework and Energy Resources of the Western Margin of Pacific Basin (Ed. The, G.H.). *Bulletin of the Geological Society of Malaysia*, 30, 195-210.
- Héroux, Y., Chagnon, A., and Bertrand, R., (1979) Compilation and correlation of major thermal maturation indicators: *American Association of Petroleum Geologists Bulletin*, 63, 2128– 2144.
- Hinz, K., Fritsch, J., Kempfer, E. H. K., Mohammad, A. M., Meyer, J., Mohamed, D., Vosberg, H., Weber, J. and Benavidez, J. (1989) Thrust tectonics along the northwestern continental margin of Sabah/Borneo. *Geologischer Rundsch.* 78, 705-730.
- Hood, A., Gutjahr, C.C.M., and Heacock, R.L., (1975) Organic metamorphism and the generation of petroleum: *American Association of Petroleum Geologists Bulletin*, 59, 986–966.
- Horne, J. C., Ferm, J. C., Carruccio, F. T., and Baganz, B. P. (1978) Depositional models in coal exploration and mine planning in Appalachian Region. In: *American Association of Petroleum Geologists Bulletin*, 62, 2379-2411.
- Hunt, J. M. (1995) petroleum geochemistry and geology, *W. H freeman and Company New York*.743p.
- Hutchison, C. S., (1989) Geological Evolution of South-East Asia. Oxford monographs on Geology and Geophysics No. 13. *Clarendon Press, Oxford*.
- James, D. M. D. (1984). The geology and hydrocarbon resources of Negara Brunei Darussalam. *Muzium Brunei*. 169 P.
- Khorasani, G. K. (1987) Novel development in fluorescence microscopy of complex organic mixtures: application in petroleum geochemistry. *Organic Geochemistry*, 11, 157-168.
- Kolaczkowska, A. E., Slougui, N. E., Watt, D. S., Marcura, R. E., and Moldowan, j. M. (1990) Thermodynamic stability of various alkylated, dealkylated and rearranged 17 α -hopane and 17 β -hopane isomers using molecular mechanic calculations. *Organic Geochemistry*, 16, 1033-1038.

References

- Koopman, A. and Schreurs, J. (1996) Onshore Lithostratigraphy. In the Geology and Hydrocarbon Resources of Negara Brunei Darussalam (Ed. Sandal, D. T.) *Brunei Shell and Brunei Museum, Bandar Seri Begawan*, 97-102.
- Kudrass, H. R., Wiedicke, M. Cepk, P. Kreuzer, H. and Muller, P. (1986) Mesozoic and Cainozoic rocks dredged from the south China Sea (Reed Bank area) and Sulu Sea and their significance for the plate-tectonic re-constructions. *Marine and Petroleum Geology*, 3, 19-30.
- Lee, C. P. (1977). The Geology of Labuan Island, Sabah, East Malaysia. BSc. Thesis University of Malaya.
- Liechti, P., Roe, R. W. and Haile, N. S. (1960) Geology of Sarawak, Brunei and Western North Borneo. Geological Survey *Department British Territories in Borneo Bulletin*. 3
- Lijmbach, G. W. M. (1975). On the origin of petroleum. Proceedings 9th world Petroleum Congress, Vol. 2, *Applied Science Publishers, London*, 357-369.
- Littke, R., Leythaeuser, D. (1993) Migration of oil and gas in coals. In: Law, B. E. Rice, D. D. (Eds.), Hydrocarbons in coal. *American Association of petroleum Geologists Studies in Geology*, 38, 219-236.
- Mackenzie, A. S., Patience, R. L., Maxwell, J. R., Vandebroucke, M., and Durand, B. (1980). Molecular parameters of maturation in the Toarcian shales, Paris Basin, France-I. Changes in the configuration of acyclic isoprenoid alkane, steranes, and triterpanes. *Geochimica et Cosmochimica Acta*, 45, 1709-1721
- Mackenzie, A. S., (1984). Application of biological markers in petroleum geochemistry. Welte, D., Brooks, J. (Eds). *Advance in petroleum geochemistry*, 1, 115-214.
- Mazlan B. Hj. Madon, (1994). The stratigraphy of northern Labuan, NW Sabah basin, East Malaysia. *Bulletin of the Geological Society of Malaysia*, 36, 19-30
- Mazlan B. Hj. Madon, (1997). Sedimentological aspects of the Temburong and Belait formations, Labuan (offshore west Sabah, Malaysia). *Bulletin of the Geological Society of Malaysia*. 41, 61-84.
- Mazlan, B. Hj. Madon, Leong Khee Meng and Azlina Anuar (1999). Sabah Bisan. In the petroleum Geology and Resources of Malaysia, 499-542. *Petroliam Nasional Berhad (PETRONAS)*.
- Moldowan, J. M., Dahl, J., Huizinga, B. J., Fago, F. J., Hickey, L. J., Peakman, T. M., Taylor, D. W., (1994) The molecular fossil recorded of oleanane and its relation to angiosperms. *Science*, 265, 768-771.
- Mukhopadhyay, P. K., Goodarzi, F., Kruege, M. A. and Alimi, M. H. (1997). Comparison of source rock geochemistry of selected rocks from the Schei Point group and Ringnes formation, Sverdrup basin, arctic Canada. *International journal of Coal Geology*, 34, 225-260.

- Murray, A. P., Sosrowidjodjo, I. B., Alexander, R., Kagi, R. I., Norgate, C. M., and Sommons, R. E., (1997). Oleanane in oil and sediments: Evidence of marine influence during early diagenesis. *Geochimica et Cosmochimica Acta*. 61, No. 6, 1261-1276.
- Ourisson, G. Albrecht P., Rohmer, M., poralla, K. (1982). Predictive microbial biochemistry from molecular fossil to prokaryotic membranes. *Trend in Biochemical Science*, 7, 236-239.
- Ourisson, G. Albrecht, P., Rohmer, M. (1984). The microbial origin of fossil fuels. *Scientific American*, 251, 44-51.
- Peakman, T. M. and Maxwell, J. R. (1988). Early diagenetic pathways of steroid alkanes. In: Advances in organic Geochemistry 1987, L. Mattavelli and L. Novelli (Eds). *Pergamon Press, Oxford*, 583-592.
- Peakman, T. M., Ten Haven, H. L., Rechka, J. R., De Leeyv, J. W., and Maxwell, J. R. (1989.) Occurrence of (22R) and (22S)- Δ 8 (H) and Δ 14 (H)- Steranes and the origin of 5α (H), 14β (H)-steranes in an immature sediment. *Geochimica et Cosmochimica Acta*. 53, 2001-2009.
- Peters, K. E. and Moldowan, J. M. (1993). In the biomarker guide interpreting molecular fossils in petroleum and ancient sediments. *Englewood Cliffs, NJ: Prentic hall*. 363.
- Philp, R. P. (1985) Fossil Fuel Biomarkers. Methods in Geochemistry and Geophysics, 23. *Elsevier, New York*, 294.
- Potter, T. L., Johans, D. R. and De Naris, T. B. G. (1984). Lithostratigraphy. In the Geology and Hydrocarbon Resources of Negara Brunei Darussalam (Ed. James, D.M.D.) *Muzium Brunei*, 43-75.
- Powell, T. G., and McKirdy, D. M. (1973). Relationship between ratio of pristane to phytane, crude oil composition and geological environment in Australia. *Nature*, 243, 37-39.
- Price, P. L., and O' Sullivan, T. and Alexander, R. (1987). The nature and occurrence of oil in Seran, Indonesia, In: Proceedings of the Indonesian Petroleum Association, 16th Annual Convention. *Indonesian petroleum Association, Jakarta*, 1, 141-173.
- Prince, R. C. (1987) Hopanoids; the world's most abundant biomolecules? *Trends in Biochemical Sciences*, 12, 455-456
- Rickard, D. T. (1970). The origin of framboids, *Lithos* 3, 269 – 293
- Riva, A., Caccialanza, P., Quagliaroli, F., (1988) Recognition of 18α (H)-oleanane in several crude and Tertiary-Upper Cretaceous sediments. *Organic Geochemistry*, 13, 671-675.
- Rullkötter, J. and Marzi, R. (1988) Natural and artificial of biological markers in a Toarcian shale from northern Germany. In: Advance in organic Geochemistry 1987, L. Mattavelli and L. Novelli (Eds.). *Pergamon Press, Oxford*, 639-645.

- Scheter, F. C. (1980). Exploration in East Malaysia over the past decade. In: Halbouy, M. T., ed., Giant Oil and Gas Fields of the Decade 1968-1978. *American Association of Petroleum Geologists Memoir*, 30, 423-440.
- Seifert W. K., and Moldowan, J. M., 1986. Use of biological marker in petroleum exploration. In: *Method in Geochemistry and Geophysics* (R. B. Johns, Ed.). 24, 261-290.
- Seifert, W. K., and Moldowan, J. M. (1979). The effect of biodegradation on steranes and terpanes in crude oils. *Geochimica et Cosmochimica Acta*. 43, 111-126.
- Seifert, W. K., and Moldowan, J. M. (1980). The effect of thermal stress on source rock quality as measured by hopane stereochemistry. *Physics and Chemistry of the Earth*, 12, 229-237
- Seifert, W. K., Moldowan, J. M. (1978). Application of steranes, terpanes, and monoaromatics to the maturation, migration, and source of crude oil. *Geochimica et Cosmochimica Acta*, 42, 77- 95.
- Shanmugam, G. (1985). Significance of coniferous rain forests and related organic matter in generation commercial quantities of oil, Gippsland Basin, Australia. *American Association of Petroleum Geologists Bulletin*, 69, 1241-1254.
- Simoneit, B.R.T. (1978). The organic chemistry of marine sediments. In: Riley, J.P., Chester, R. (Eds.), *Chemical Oceanography 7*. Academic Press, New York, 233-311.
- Stach, E., Mackowsky, M. T., Teichmuler, M., Taylor, G. H., Chandra, D., Teichmuler, R. (1982). Stach's Textbook of Coal Petrology, 3edn. *Gebrüder Borntraeger, Berlin*, 535 p.
- Stopes, M. C. (1935). On the petrology of banded bituminous coals. *Fuel*, 14, London.
- Tan, D. N. K. and Lamy, J. M. (1990) Tectonic evolution of the NW Sabah continental margin since the late Eocene. *Bulletin of the Geological Society of Malaysia*, 27, 241-260
- Taylor, G. H., Teichmüller, M., Davis, A., Diessel, C. F. K. (1998). Organic Petrology: A new handbook incorporating some revised parts of Stach's text book of Coal Petrology. *Gebrüder Borntraeger, Berlin-Stuugart*.
- Teichmuller, M. (1974). Generation of petroleum-like substances in coal seams as seen under the microscope. In: Tissot, B. and Bienner, F. (Eds.), *Advances in Organic geochemistry 1973*. *Editions Technip, Paris*, 321-348.
- Teichmuller, M. (1989). The genesis of coal from the viewpoint of coal petrology. *International Journal of Coal Geology*, 12, 1-87.
- Tissot, R. P., Welte, D. H. (1984). Petroleum formation and occurrence. 2nd edition, *Springer, Berlin Heidelberg New York*, 699 p.

- Tongkul, F. (2001). Sumber geologi intrinsik Pulau Labuan. Warisan Geologi Malaysia, 4, 377- 387
- Volkman, J. K., and Maxwell, J. R. (1986). Acyclic isoprenoids as biological markers. In: Biological Markers in the Sedimentary Record (R. B. Johns, Ed.) Elsevier, New York, 1-42.
- Wan Hasiah Abdullah (2001). A petrographic comparison of oil-generating coals from the tropics and non oil-generating coals from the arctic, *Geol. Soc. Malaysia Ann. Geol. Conf. 2001*, June 2-3 2001, Pangkor Island, Perak Darul Ridzuam, Malaysia, p. 33-38
- Wan Hasiah Abdullah (1997a). Evidence of early generation of liquid hydrocarbon from suberinitic as visible under the microscope, *Organic Geochemistry*, 27, 591-596
- Wan Hasiah, Abdullah (1997b). Common liptinitic constituents of Tertiary coals from the Bintulu area and Merit-Pila coalfields, Sarawak and their relation to oil generation from coal. *Bulletin of the Geological Society of Malaysia*, 41, 85-94
- Wan Hasiah, Abdullah (1999a). Oil-generating potential of Tertiary coals and other organic rich sediments of Nyalau Formation, onshore Sarawak. *Journal of Asian Earth Sciences*, 17, 225-267.
- Wan Hasiah, Abdullah (1999b) Organic facies variations in the Triassic shallow marine and deep marine shales of central Spits Bergen, Svalbard, *Marine and Petroleum Geology*, 16, 467-481.
- Waples, D. W. and Machihara, T. (1990). Application of sterane and triterpane biomarkers in petroleum exploration. *Bulletin of Canadian Petroleum Geology*, 38, No. 3, 357-380.
- Wilkins, R. W. T., George, S. C. (2002), Coal as source rock for oil: a review, *International journal of Coal Geology*, 50, 317-361.
- Wilson, R. A. M. (1964). The Geology and Mineral Resource of the Labuan and Padas Valley Area, Sabah, Malaysia. *Geological Survey Borneo Region, Malaysia, Memoir 17, 150p*
- Wood, B. G. M. (1985). The mechanics of progressive deformation in crustal Plate- a working model for Southeast Asia. *Bulletin of the Geological Society of Malaysia*, 18, 55-99.
- Zumberge, J. E. (1984). source rocks of the La Luna Formation (Upper Cretaceous) in the middle Magdalena Valley, Colombia. In: *Geochemistry and Source Rock Potential of Carbonate Rocks*, J. G. palacas (ed.) AAPG Studies in Geology No. 18. *American Association of Petroleum Geologists*, 127-133.